PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

A61K 48/00, C07H 21/04

A1

(11) International Publication Number: WO 99/55380

(43) International Publication Date: 4 November 1999 (04.11.99)

(21) International Application Number: PCT/US99/09039
 (22) International Filing Date: 27 April 1999 (27.04.99)

(30) Priority Data:

60/083,152 27 April 1998 (27.04.98) US 60/112,098 14 December 1998 (14.12.98) US

(71) Applicant: PACIFIC NORTHWEST CANCER FOUNDA-TION [US/US]; 120 Northgate Plaza, Seattle, WA 98125 (US).

(72) Inventors: MURPHY, Gerald, P.; 1134 23rd Avenue E., Seattle, WA 98112 (US). BOYNTON, Alton, L.; 2420 274th Avenue N.E., Redmond, WA 98053 (US). SEHGAL, Anil; Apartment H106, 2104 North 105th Street, Seattle, WA 98133 (US).

(74) Agents: BALDWIN, Geraldine, F. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: Nr-CAM GENE, NUCLEIC ACIDS AND NUCLEIC ACID PRODUCTS FOR THERAPEUTIC AND DIAGNOSTIC USES FOR TUMORS

(57) Abstract

The present invention relates to the identification of a novel role of Nr-CAM in cell transformation and aberrant cellular proliferation. In particular, the present invention relates to the altered gene expression of Nr-CAM in a number of primary tumors and cell lines derived from tumors, in addition to, the altered gene expression of ligands for Nr-CAM. Further, the present invention relates, in part, to the Applicants' surprising discovery that the inhibition of Nr-CAM gene expression or the inhibition of -CAM activity in transformed cells reverses the transformed phenotype.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Albania	ES	Spain	LS	Lesotho	SI	Slovenia	ì
AL AM	Amenia	FI	Pinland	LT	Lithuania	SK	Slovakia	
AM	Amenia	FR	France	LU	Luxembourg	SN	Senegal	
		GA	Gabon	LV	Latvia	SZ	Swaziland	
AU -	Australia	GA GB	United Kingdom	MC	Monaco	TD	Chad	
AZ	Azerbaijan	GB GE	Georgia	MD	Republic of Moldova	TG	Togo	
BA	Bosnia and Herzegovina	GE GH	Ghana	MG	Madagascar	TJ	Tajikistan	
BB	Barbados	GH GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan	
BE	Belgium	GN GR	Guinea	MIK	Republic of Macedonia	TR	Turkey	
BF	Burkina Faso	-		ML	Mali	TT	Trinidad and Tobago	
BG	Bulgaria	HU	Hungary	MN		UA	Ukrainė	
BJ	Benin	IE	Ireland		Mongolia	UG	•	
BR	Brazil	IL	Israel	MR	Mauritania		Uganda	
BY	Belarus	IS .	Iceland	MW	Malawi	US	United States of America	
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan	
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam	
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia	
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe	
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand			
CM	Cameroon		Republic of Korea	PL	Poland			_
CN	China	KR	Republic of Korea	PT	Portugal			
CU	Cuba	KZ	Kazakstan	RO	Romania			
cz	Czech Republic	LC	Saint Lucia	RU	Russian Federation			
DE	Germany	LI	Liechtenstein	SD	Sudan			
DK	Denmark	LK	Sri Lanka	SE	Sweden			
RE	Estonia	LR	Liberia	SG	Singapore			
					•			
i								

Nr-CAM GENE, NUCLEIC ACIDS AND NUCLEIC ACID PRODUCTS FOR THERAPEUTIC AND DIAGNOSTIC USES FOR TUMORS

The present application is entitled to and claims priority benefits of U.S. Application No. 60/083,152, filed April 27, 1998; and 60/112,098, filed December 14, 1998, the entire disclosures of which are incorporated herein by reference.

1. FIELD OF THE INVENTION

The present invention relates to the identification of a novel role for the neuron-glia-related cell adhesion molecule (Nr-CAM) gene in tumorigenesis, in particular primary brain tumorigenesis. The present invention is related to the role of Nr-CAM nucleic acids and polypeptides as diagnostic tools to indicate a pre-cancerous condition or cancer, and therapeutic agents based thereon to inhibit Nr-CAM gene expression and/or activity as a method of treating, inhibiting and/or preventing tumorigenesis.

2. BACKGROUND OF THE INVENTION

20

2.1. BRAIN TUMORS

Brain tumors are among the leading cause of death among young children and adults. A survey by the American Cancer Society has documented that 13,300 people died of brain tumors in 1995 and predicated that over 17,900 would die in 1996 (Parker et al., 1996, CA Cancer J. Clin., 46:5-28). The number of deaths due to brain tumors has been increasing at a significant rate each year. On average, 25,000 Americans are diagnosed with brain cancer yearly. Brain tumors claim the lives of more children than any other form of cancer except leukemia.

The increased incidence of brain tumors is not only evident in children but also in adults. It has been documented that a significant increase in mortality has

occurred in adult primary malignant tumors between 1982 and 1996 (Parker et al., 1996). Glioblastomas, astrocytomas and meningiomas are the most common brain tumors that affect adults (Thapar and Laws, 1993, CA Cancer J. Clin., 43:263-₅ 271).

The transformation of normal human brain cells into gliomas occurs as a result of the accumulation of a series of cellular and genetic changes (Sehgal, 1998, Cancer Surv., 25:233-275; vonDiemiling et al., 1995, Glia 15:328-338; Furnari et al., 1995, J. Surg. Oncol. <u>67</u>:234). 10 alterations include the loss, gain or amplification of different chromosomes. These genetic changes lead to altered expression of proteins that play important roles in the regulation of normal cell proliferation. Several common genetic alternations at the chromosomal level (loss of 17p, 15 13q, 9p, 19, 10, 22q, 18q and amplification of 7 and 12q) have been observed (Sehgal et al., 1998 J. Surg. Oncol. 67:23; vonDiemiling et al., 1995, Glia 15:328-338; Furnari et al., 1995, Cancer Surv. 25:233-275). These alterations lead to changes in the expression of several genes (p53, RB, 20 INFα/β, CDKN2, MMAC1, DCC, EGFR, PDGF, PDGFr, MDM2, GLI, CDK4 and SAS) during the genesis and progression of human gliomas (Sehgal, 1998, J. Surg. Oncol. 67:234; vonDiemiling et al., 1995, Glia 15:328-338). Recent studies have suggested that altered expression of several other genes (MET, MYC, TGF\$, CD44, VEGF, N-CAML1, p21waf1/Cip1, trkA, MMRs, C4-2, D2-2) and proteins (cathepsins, tenascin, matrix metalloproteases, tissue inhibitors of metalloproteases, nitric oxide synthetase, integrins, IL 13 receptor, Connexin 43, uPAR's extracellular matrix proteins and heat shock proteins) are associated with the genesis of human gliomas (Sehgal, 1998, J. Surg. Oncol. 67:234). Taken together these findings point

to the fact that accumulation of multiple genetic mutations coupled with extensive changes in gene expression may be a

prerequisite in the etiology of human gliomas. Despite identification of these genetic alterations, the exact series of events that leads to the genesis of human gliomas is not clear.

Glioblastoma multiforme are high grade astrocytomas 5 that grow very rapidly and contain cells that are very malignant (Thapar and Laws, 1993, CA Cancer J. Clin., 43:263-The molecular basis of glioblastoma multiforme occurrence may involve systematic events at the chromosomal level or at a gene expression level. These may include inactivation of tumor suppressor genes, activation of oncogenes or specific translocations at the chromosomal level. Some genetic changes at the chromosomal level and gene expression level have been well documented for other brain tumors (Furnari et al., 1995, Cancer Surv., 25:233-**15** 275). For example, it has been documented that loss of tumor suppressor(s) genes at chromosome 10, mutations in p53, or overexpression of epidermal growth factor receptor, may be major events leading to glioblastoma multiforme. A number of other genes such as EGFR, CD44, \$4 integrins, membrane-type 20 metalloproteinase (MT-MMP), p21, p16, p15, myc, and VEGF have been shown to be overexpressed in different types of brain tumors (Faillot et al., 1996, Neurosurgery, 39:478-483; Eibl et al., 1995, J. Neurooncol., 26:165-170; Previtali et al., 1996, Neuropathol. Exp. Neurol. <u>55</u>:456-465; Yamamoto et al., 1996, Cancer Res., <u>56</u>:384-392; Jung et al., 1995, Oncogene, 11:2021-2028; Tsuzuki et al., 1996, Cancer, 78:287-293; Chen et al., 1995, Nature Med., 1:638-643; Takano, et al., 1996, Cancer Res., 56:2185-2190; Bogler et al., 1995, Glia, 15:308-327). Several cell adhesion molecules (CAMs), such as integrins, cadherins, IgSF proteins (carcinoembryonic antigen, N-CAM and VCAM-1) or lectins, are thought to be involved in tumorigenesis (Johnson, 1991, Cancer Metastat. Rev. 10:11-22). Over-expression of anti-sense to the

secreted glycoprotein SPRAC (secreted protein, acidic and rich in cysteine), results in suppression of the adhesive and invasive capacities of melanomas (Ledda et al., 1997, Nature (Med). 3:171-176). The cell-surface adhesion molecule 5 MCAM(MUC18) when over-expressed may lead to increased adhesion and metastatic potential of human melanoma cells in nude mice (Xie et al., 1997, Proc. Nat'l Cancer Conf. 38:522). Expression of N-CAM or ICAM (Intracellular adhesion molecule) is related inversely to increased metastasis (Hortsch, 1996, Neuron 17:587-593). Other genes such as p53 10 show mutations in the majority of brain tumors (Bogler et al., supra). How the interplay of one or more of these genes leads to tumorigenesis is not known but most likely multiple steps are required for neoplastic transformation. The exact series of events that lead to initiation or progression of 15 glioblastoma are not known at present and useful markers for early detection of brain tumors are lacking.

2.2. <u>CAMB</u>

A subfamily of the immunoglobulin superfamily
20 (IgSF) proteins are termed "cell adhesion molecules" (CAMs)
(Hortsch, 1996, Neuron 17:587-593).

Several CAM family members are implicated in the process of tumorigenesis including, N-CAM, CEA, (Cacinoembryonic Antigen), DCC (Deleted in Colon Carcinoma) and L1.

CEA is a cell surface glycoprotein of colon mucosal cells. High levels of CEA are observed in the serum of tumor patients (Benchimol, et al., 1989, Cell, <u>57</u>:327-334). It was demonstrated that over-expression of CEA in malignant cells may disturb intercellular adhesion that may in turn cause tissue disruption leading to the metastasis of primary tumor cells (Albelda, 1993, Lab Invest, <u>68</u>:4-14; Benchimol et al., 1989).

L1 is another cell adhesion molecule that belongs to the Ig superfamily and is expressed in neuroblastomas, melanomas, lymphocytes and Schwann cells (Izumoto, et al., 1996, Cancer Res, 56:1440-1444). Antibody neutralization experiments demonstrate that L1 is responsible for the highly invasive nature of the C6 glioma cells. Mutations in the L1 are known to be associated with a spectrum of neurological deficiencies including mental retardation (Izumoto et al., 1996).

The neural cell adhesion molecules (N-CAMs) are predominantly though not exclusively, expressed in developing peripheral and central nervous systems of a number of invertebrates and vertebrates. These proteins are generally present on the cell surface and consist of multiple Ig domains, multiple fibronectin type III repeats near the cell membrane and either a transmembrane domain or a glycophosphatidylinositol-linked membrane anchor at the C-terminus (Hortsch, 1996). The N-CAMs can be grouped into 2 major structural families, one resembling N-CAM and the other resembling the liver CAM (L-CAM) and its mammalian homologue uvomorulin or E-cadherins. Within N-CAM, 2 major types are observed, the N-CAM (neuronal CAM) and the neuron-glial CAM (Ng-CAM) (Grumet et al., 1991, J. Cell Biol. 113:1399-1412).

N-CAM is expressed in a wide variety of tissues and is implicated in embryonic development. Over-expression of N-CAM is observed in a variety of tumors including multiple myelomas, small cell carcinomas and adenoid cystic carcinomas. Down regulation of N-CAM is observed in malignant glioma cells (Albelda, 1993; Poley, et al., 1997, Anticancer Research, 17:3021-3024). In Wilm's tumor of kidney, the N-CAM exists in h-PSA form that is less adhesive to surrounding cells and fibers. In a recent study it was demonstrated that high levels of N-CAM were detected in patients with prostate carcinomas (Lynch, et al., 1997,

Prostate, 32:214-220). In vivo studies in nude mice demonstrated that N-CAM may be involved not only in adhesive and motile behavior of cells but also in their growth regulation (Poley et al., 1997; Lynch et al., 1997).

DCC is a cell adhesion molecule that belongs to the N-CAM family. DCC was first shown to be expressed in a variety of tumors including the brain and lung but its expression was reduced and mutated in a number of colorectal carcinomas (Fearon, et al., 1990, Cell, 61:759-767). The down regulation or mutation of the DCC molecule lead to the disruption of normal cell-cell adhesion in the intestinal epithelium. This process is known to play an important role in the metastasis of colorectal carcinomas (Albelda, 1993; Fearon et al., 1990).

15

2.2.1. Nr-CAM

Nr-CAM (neuron-glia related CAM) was cloned from a chicken brain library when Ng-CAM cDNA Clones were being isolated (Grumet et al., 1991, J. Cell Biol. <u>113</u>:1399-1412). Monoclonal antibodies against E8 tectal surface protein 20 identified a similar molecule that was cloned from a chicken brain library. This protein was designated Bravo/Nr-CAM (Grumet et al., 1991; Lane et al., 1986, Genomics 35:456-465). The Nr-CAM protein contains 6 Ig domains and 5 fibronectin repeats. Numerous studies on chicken Nr-CAM suggested that it may play an important role in cell-cell adhesion during the development of the vertebrate nervous system. The human homologue of the chicken Nr-CAM has been cloned (Lane et al., 1996, Genomics 35:456-465; see Lane et al., Figure 1 at pages 458-9 for nucleotide and deduced amino acid sequences of hNr-CAM, incorporated herein by reference). 30

Sequence analysis of Nr-CAM proteins isolated from human rat, chicken, and mouse showed more than 80% identity. One unique characteristic of hNr-CAM is that the third

fibronectin repeat contains a furin-like cleavage site (Grumet, 1991, Current Opinion in Neurobiology, 1:370-376; Suter, et al., 1995, J. Cell Biol., 131:1067-1081). It has also been reported that the 140KDa protein may exist as a doublet (Grumet, et al., 1991, J. Cell Biol., 113:1399-1412; Lane, et al., 1996, Genomics, 35:456-465).

The cytoplasmic tail of the hNr-CAM protein is known to interact with a cytoplasmic protein named ankyrin (Davis, et al., 1993, J. Cell Biol., 121:121-133; Davis, et al., 1996, J. Cell Biol., 135:1355-1367). This region of the hNr-CAM protein is highly conserved among other family members (Ng-CAM, L1, neurofascin and neuroglian protein). Neurofascin and L1 proteins contain a phosphorylation site that may modulate its interaction with ankyrin (Davis, 1993; Davis 1996). Phosphorylation status of hNr-CAM in this region of the protein has not been reported yet. The cytoplasmic tail of the hNr-CAM contains sequences that have potential to interact with PSD-95/discs-large/ZO-1 family of membrane associated proteins (Grumet, et al., 1991, J Cell Biol, 113:1399-1412).

- cDNA analysis of rat Nr-CAM revealed two different forms of Nr-CAM as a result of alternative mRNA splicing. The alternatively spliced form of Nr-CAM contains 10-amino acids inserted between the Ig domain and fibronectin domain, and 15-amino-acids inserted after the fourth fibronectin repeat, and complete deletion of the fifth fibronectin repeat (Suter et al., 1995). The extracellular portion of the Nr-CAM protein contains twenty potential sites for N-linked glycosylation (Kayyem, et al., 1992, J Cell Biol, 118:1259-1270; Suter et al., 1995; Grumet, 1991; Cell Tissue Res.

 290:423-428).
- Nr-CAM is expressed in growing neurites and radial cells of the optic chiasm. Nr-CAM protein family members not only play an important role in cell adhesion but they can

interact with other proteins, such as FGF-R, by phosphorylation events to bring about neurite extension. L1-CAM gene mutation may be involved in several neurological disorders (Hortsch, 1996). A number of transformed cells from a variety of tissues also express L1-CAM, and the expression is correlated inversely with the metastatic capacity of a lymphoma cell line in mice, leading to speculation that these CAMs may play a role in metastatic events (Hortsch, 1996). See generally, Albelda, 1993, Lab. Invest. 68:4-14.

It has been demonstrated that hNr-CAM is a brain specific protein and is expressed on neurons, Schwann cells, Muller cells and transiently in the cells of floor plate (Davis, et al., 1996, J Cell Biol, 135:1355-1367; Grumet, Cell Tissue Res, [1991] 290:423-428). The hNr-CAM protein is expressed preferentially on fiber tracts, in spinal cord, cerebellum, tectum, and telencephalon. It is also known to be concentrated in the node of Ranvier, thus demonstrating its potential role in the formation and maintenance of the nodes (Suter, et al., 1995, J Cell Biol, 131:1067-1081). The hNr-CAM is also expressed widely in the retina on cell bodies and fiber layers and on the ganglion cells (Suter et al., 1995; Davis, et al., 1994, J Biol Chem, 269:27163-27166).

The hNr-CAM protein is a cell adhesion molecule that can function as a receptor and as a ligand. It not only interacts with other hNr-CAM molecules on the cell surface but also with other CAMs and extracellular matrix proteins. Some of the proteins that interact with the hNr-CAM include: contactin/F11, axonin-1 ax-1), neurofascin (Nf), Receptor protein tyrosine phosphataseβ (RPTPβ), Ng-CAM, chondroitin sulfate proteoglycans neurocan, and phosphacan (Davis et al., 1994; Grumet et al., 1991; Cell Tissue Res. 290:423-428).

Recent analysis has indicated that the cytoplasmic domain of hNr-CAM can directly interact with ankyrins, a family of

spectrin-binding proteins (Davis, 1993; Davis, 1996). This interaction may be involved in sending signals from the extracellular domain to the cytoplasm of cells. Antibody binding analysis during the tectal neurite growth has demonstrated that the hNr-CAM can act as a receptor and as a ligand. Some of the major functions of the hNr-CAM are: a) modulation of axonal growth and guidance; b) modulation of the function of non-neuronal glial cells; c) synapse formation and maturation of the central nervous system; and d) as a heterophilic neuronal receptor (Grumet, 1991 Cell Tissue Res. 290:423-428).

Citation of references herein shall not be construed as an admission that such references are prior art to the present invention.

3. <u>SUMMARY OF THE INVENTION</u>

The present invention relates to the discovery of a novel role for Nr-CAM in the aberrant proliferative behavior of a number of cell types, including numerous primary tumors and derived cell lines. In particular, the present invention 20 relates to the identification of the role of Nr-CAM in cell transformation and tumorigenesis, in particular, brain. present invention encompasses therapeutic and diagnostic applications based on Nr-CAM proteins, nucleic acids, and agonists and antagonists, for the treatment, inhibition or prevention of tumorigenesis. The present invention further encompasses therapeutic and diagnostic applications based on a ligand of Nr-CAM proteins, nucleic acids, and agonists and antagonists, for the treatment, inhibition or prevention of tumorigenesis. The present invention further encompasses screening assays to identify modulators of Nr-CAM activity 30 and/or expression as potential therapeutic agents for the treatment, inhibition and/or prevention of a transformed phenotype or tumorigenesis.

The present invention is based, in part, on the Applicants' surprising discovery that the Nr-CAM nucleotide sequence and encoded protein product is expressed at high levels in glioblastoma multiforme tissue, astrocytomas, gliomas, glioblastoma tumor tissues, as well as, certain other forms of tumors and cancers.

In one embodiment, the present invention encompasses nucleotide sequences complementary to the nucleotide sequence of Nr-CAM, such as primers, fragments or antisense nucleotides which may be used to determine the level of Nr-CAM expression in a tissue or cell culture sample as prognostic of a pre-cancerous or transformed cell phenotype; or to inhibit Nr-CAM expression as a method of treating, inhibiting or preventing a pre-cancerous or transformed cell phenotype. In a specific embodiment, the Nr-CAM gene is a human gene and the Nr-CAM protein is a human protein.

The present invention also encompasses inhibitors of Nr-CAM activities related to cellular transformation. Nr-CAM is a known protein thought to play a role in cell-cell 20 adhesion, e.g. during development of not only the vertebrate nervous system but also by interaction with other proteins, such as FGF-R, in neurite extension. The present invention encompasses peptide fragments or antagonists, antibodies, or small compounds which may inhibit or compete with ligands binding to Nr-CAM and thus inhibit Nr-CAM activity. 25 invention further encompasses peptide fragments (and derivatives and analogs thereof) which comprise one or more domains of a Nr-CAM protein which may be used to prevent ligands binding to Nr-CAM. Antibodies to Nr-CAM, and to Nr-CAM derivatives and analogs, are additionally provided. Methods of production of the Nr-CAM proteins, derivatives and

analogs, e.g., by recombinant means, are also provided.

The present invention further encompasses screening assays to identify compounds which inhibit Nr-CAM nucleic acid expression or nucleic acid product activity as potential therapeutics for the treatment and/or prevention of tumorigenesis. In particular, the present invention encompasses host cell lines or transgenic animals which express Nr-CAM at high levels which have utility as tools for screening assays to identify agents which inhibit Nr-CAM expression and/or activity as potential therapeutic agents for the treatment and prevention of tumorigenesis.

The present invention also encompasses therapeutic and diagnostic methods and compositions based on Nr-CAM proteins and nucleic acids. Therapeutic compounds of the invention include but are not limited to Nr-CAM proteins and analogs and derivatives (including fragments) thereof; antibodies thereto; nucleic acids encoding the Nr-CAM proteins, analogs, or derivatives; and Nr-CAM antisense nucleic acids.

10

The invention provides for treatment of disorders of overproliferation (e.g., tumors, cancer and 20 hyperproliferative disorders) by administering compounds that decrease or antagonize (inhibit) Nr-CAM function (e.g., antibodies, antisense nucleic acids, ribozymes).

The invention also provides methods of treatment of disorders involving deficient cell proliferation (growth) or in which cell proliferation is otherwise desired (e.g., degenerative disorders, growth deficiencies, lesions, physical trauma) by administering compounds that promote Nr-CAM activity (e.g., an agonist of Nr-CAM; nucleic acids that encode Nr-CAM).

Animal models, diagnostic methods and screening methods for predisposition to disorders, and methods for identification of Nr-CAM agonists and antagonists, are also provided by the invention.

3.1. DEFINITIONS AND ABBREVIATIONS

As used herein, underscoring or italicizing the name of a gene shall indicate the gene, in contrast to its encoded protein product, which is indicated by the name of the gene in the absence of any underscoring or italicizing. For example, "Nr-CAM" shall mean the Nr-CAM gene, whereas "Nr-CAM" shall indicate the protein product of the Nr-CAM gene.

As used herein, the following terms shall have the meanings indicated.

Nr-CAM nucleotides or coding sequences: DNA sequences encoding Nr-CAM mRNA transcripts, protein, polypeptide or peptide fragments of Nr-CAM protein, and Nr-CAM fusion proteins, and RNA sequences corresponding to Nr-CAM mRNA transcripts and RNA sequences which are

complementary to the mRNA transcript, Nr-CAM nucleotide sequences encompass RNA, DNA, including genomic DNA (e.g. the Nr-CAM gene) and cDNA.

Nr-CAM: gene products, e.g., transcripts and the Nr-CAM protein. Polypeptides or peptide fragments of the 20 protein are referred to as Nr-CAM polypeptides or Nr-CAM peptides. Fusions of Nr-CAM protein, polypeptides, or peptide fragments to an unrelated protein are referred to herein as Nr-CAM fusion proteins.

As used herein, the following terms shall have the abbreviations indicated.

CD: cytoplasmic domain

DD-PCR: differential display - polymerase chain

reaction

ECD: extracellular domain

30 FNHA: fetal_normal_human_astrocytes

GMTT: glioblastomas multiforme tumor tissue

MTB: multiple tissue blot

MTT: meningioma tumor tissue

NBT: normal brain tissue ORF: open reading frame

RT-PCR: reverse transcription - polymerase chain

5 reaction

TM: transmembrane domain UTR: untranslated region

Brain tumor cell lines:

10 CCF-STTG1: astrocytoma grade IV

D283 Med: medulloblastoma

DBTRG-05MG: glioblastoma multiforme

Hs 683: glioma

IMR-32: neuroblastoma

PFSK-1: primitive neuroectodermal tumor

SW 1783: astrocytoma grade III

4. DESCRIPTION OF THE FIGURES

Figure 1 is an autoradiogram of DD-PCR gel illustrating expression of hNr-CAM (designated D4-1) and of a control D1-2 gene in normal brain tissue (N) and brain tumor (T) tissue, i.e., glioblastoma multiforme (GM).

Figures 2 (A-D) present the nucleotide and amino acid sequences of human Nr-CAM has well as the results of nucleotide sequence analysis as described in Section 6 (Figure 2C) and a schematic illustration of the hNr-CAM gene showing the area used herein for antisense targeting (Figure 2D). Figure 2A presents the nucleotide sequence of human Nr-CAM. Features of the nucleotide sequence include the following: Nucleotides 130-3615 encode the extracellular domain; nucleotides 202-4026 encode product = hBRAVO-Nr-CAM; nucleotides 316-483 encode the Immunoglobulin I domain; nucleotides 613-768 encode the Immunoglobulin II domain; nucleotides 988-1134 encode the Immunoglobulin III domain;

nucleotides 1258-1410 encode the Immunoglobulin IV domain; nucleotides 1540-1719 encode the Immunoglobulin VI domain; nucleotides 2113-2265 encode the first Fibronectin (Fn) repeat; nucleotides 2413-2565, the second Fn repeat; nucleotides 2710-2886, the third Fn repeat; nucleotides 3028-3186 the fourth Fn repeat; nucleotides 3370-3510, the fifth Fn repeat; nucleotides 2616-3684, the transmembrane region; nucleotides 3685-4036, the intracellular domain; and nucleotides 4030-4134 constitute a 3' untranslated region. Figure 2B presents the amino acid sequence of human Nr-CAM.

The hydrophobic signal sequence is underlined. Figures 2A and 2B are adapted from Lane et al., 1996, Genomics 35:456-465.

Figure 2C illustrates nucleotide sequence identity analysis between previously cloned hNr-CAM (Accession Number U55258), rat Nr-CAM (Accession Number U81037) and the sequence of clone D4-1 obtained by cloning the hNr-CAM isolated by DD-PCR into pCRII vector (Invitrogen). Sequence identity analysis was performed using the DNasis program from Hitachi Software (South San Francisco, CA). Stars (*) indicate presence of identical nucleotides among the sequences.

Figure 2D presents a schematic of the hNr-CAM gene, including, in particular, the area used herein for antisense targeting. The arrow indicates this area. See text, Section 7 for details.

25

Figures 3 (A-F) illustrate differential expression of Nr-CAM in glioblastoma multiforme tissue (GM) when compared to normal brain tissue (NT). The technique of in situ hybridization was used. Figures 3 (A, B and C) are from one GM tumor while Figures 3 (D, E and F) are from a second GM tumor. Figures 3 (A and D) show tumor regions and one hybridized with hNr-CAM anti-sense probes. Figures 3 (B and E) show serial sections of A and D, and are hybridized with

hNr-CAM sense probes. Figures 3 (C and F) show normal brain
regions of serial sections of the same brain as sections A
and D, respectively, and are hybridized with hNr-CAM antisense probe. Cells expressing hNr-CAM are indicated by
arrows.

Figures 4 (A-B) illustrate differential expression of hNr-CAM in normal and tumor tissues. Figure 4A is a gel obtained using RT-PCR of total RNA of a variety of tissues. The upper panel of Figure 4A shows expression of hNr-CAM and the lower panel shows expression of a central, housekeeping gene D1-2. Figure 4B is a bar graph showing relative expression of hNr-CAM after correction for gel load based on D1-2 expression. In both Figures 4A and 4B, the specific tissue in each of lanes 1-14 is as follows: Lane 1, glioblastoma IV; 2, recurrent meningioma; 3, meningioma; 4, normal brain (GS); 5, neuroblastoma; 6, recurrent glioma; 7, glioblastoma multiforme; 8, melanoma; 9, normal breast; 10, tumor breast; 11, benign prostate; 12, prostate tumor; 13, normal brain; and 14, glioma III. Recurrent gliomas represent a tissue from the same patient previously diagnosed 20 with GM.

Figures 5 (A and B) illustrate differential expression of hNR-CAM in normal brain tissue and astrocytoma tumor tissue. Figure 5A is a gel obtained using RT-PCR of total RNA of normal or astrocytoma tissue. The upper panel of Figure 5A shows expression of hNr-CAM; the middle panel, EGFR (epidermal growth factor receptor, a known brain tumor maker) and the lower panel, D1-2. Figure 5B is a bar graph showing the relative expression of hNr-CAM and EGFR after normalizing with D1-2 gel loading. In both Figures 5A and 5B, Lane a is normal brain tissue and Lane b is astrocytoma tumor tissue.

Figures 6 (A and B) illustrate expression of hNr- CAM in normal brain and brain tumor cell lines. Figure 6A is

an autoradiogram of a Southern blot of hNr-CAM expression (upper panel); and of D1-2 expression (lower panel) in various brain cell lines. Figure 6B is a bar graph showing relative expression of hNr-CAM in respective cell lines after correction for gel loading based on D1-2 expression. In both Figures 6A and 6B, the specific cell line in each of Lanes 1-9 is as follows: Lane 1, astrocytoma III; 2, astrocytoma IV; 3, glioblastoma; 4, glioma; 5, neuroectodermal; 6, medulloblastoma; 7 neuroblastoma; 8, FNHA; and 9, normal brain.

Figures 7 (A and B) illustrate expression of hNr-CAM in different regions of the brain. Figure 7A is an autoradiogram of a Northern blot of hNr-CAM expression (upper panel) and of β-actin expression (lower panel) in a variety of regions (Lanes 1-8) of normal adult brain. β-actin expression serves as an internal control for gel loading for different regions of the brain. Figure 7B is a bar graph showing the relative expression of hNr-CAM in different brain regions after correction for gel loading based on β-actin expression. In both Figures 7A and 7B, the specific region of brain in each of Lanes 1-8 is as follows: Lane 1, cerebellum; 2, cerebral cortex; 3, medulla; 4, spinal cord; 5, occipital pole; 6, frontal lobe; 7, temporal lobe; and 8, putamen.

Figures 8 (A and B) illustrate expression of hNr
CAM in human cancer cell lines (Lanes a-h). Figure 8A is an autoradiogram of a Northern blot of hNr-CAM expression (upper panel) and of β-actin expression (lower panel). Figure 8B is a bar graph of the relative expression of hNr-CAM after correction for gel loading based on β-actin expression. In both Figures 8A and 8B, the specific human cancer cell line in each of Lanes a-h is as follows: Lane a, promyelocytic leukemia (HL-60); b, HeLa cells (S3); c, chronic myelogenous leukemia (K-562); d, lymphoblastic leukemia (MOLT-4); e,

Burkitt's lymphoma (Raji); f, colorectal adenocarcinoma (SA 480); g, lung carcinoma (A549); and h, melanoma (G361).

Figure 9 illustrates genomic Southern blot analysis of hNr-CAM in several brain tumor cell lines and in the NIH 3T3 cell line. Upper panel shows an ethidium bromide-stained gel of genomic DNA digested with EcoRI restriction enzyme. Lower panel shows an autoradiogram of the Southern blot. The arrow indicates hNr-CAM. In both panels, the brain tumor cell lines in each of Lanes 1-4 is as follows: Lane 1, NIH 3T3 cell line; 2, astrocytoma III; 3, glioma and 4, glioblastoma cell line.

Figures 10 (A and B) illustrate the effects of antisense hNr-CAM expression on the morphology of glioblastoma (GB) cells. Figure 10A shows GB cells transfected with p-CMV-neovector only. Figure 10B shows GB cells transfected with p-CMV-neovector containing antisense hNr-CAM.

Figure 11 is a graph illustrating the effect of anti-sense hNr-CAM expression on the proliferation of glioblastoma cells.

Figure 12 is a bar graph illustrating the effect of anti-sense hNr-CAM on soft agar colony formation of glioblastoma cells. GB represents Glioblastoma cells; GB-PFCS, represents glioblastoma cells with vector alone; GB-Anti-Nr-CAM, represents GB cells expressing Nr-CAM 1/3.

Figure 13 illustrates the effect of overexpression of antisense hNr-CAM (pCMV-1/3 Nr-AS) on native hNr-CAM mRNA level. See text Section 7.1.6. for details.

Figures 14 (A and B) schematically illustrate the effect of antisense hNr-CAM expression in glioblastoma cells. Figure 14A shows untransfected cells and Figure 14B shows

9CMV-1/3Nr-AS transfected cells.

Figures 15 (A-D) illustrate effect of antisense hNr-CAM expression on morphology of glioblastoma (GB) cells.

Figures 15 (A and B) show 5GB cells transfected with pCMVneovector only, i.e., control cells (at different
magnifications). Figures 15 (C and D) show 5 GB cells
transfected with pCMV-1/3Nr-AS (at different magnifications).

5 Spindle shaped cells are indicated by arrows. See text
Section 7.2.1. for details.

Figures 16 (A-D) show the effect of serum treatment on the morphology of the pCMV-1/3Nr-AS transfected 5GB cells.

5GB (pCMVneo or pCMV-Nr-1/3AS transfected) cells were plated at an approximate density of 3x10⁴. Both cell types were treated with different concentrations of FBS. Figures 15 A, B, C, and D, respectively, show 5GB cells (pCMV-1/3Nr-CAM transfected) treated with 0.1, 1.0, 2.0 and 5% FBS serum.

Figure 17 shows the effect of antisense hNr-CAM pCMV-1/3Nr-AS) expression on the proliferation of 5GB human qlioblastoma cell line.

Figure 18 shows the effect of anti-sense hNr-CAM overexpression on the migration ability of 5GB cells.

Briefly, 1x10⁶ pCMV-neo and pCMV-1/3Nr-AS transfected cells were plated in triplicates in transwell inserts (8μm) from 20 Coastar (Cambridge, MA). Inserts were placed in 6 well tissue culture dishes containing the appropriate growth medium with serum. Cells that migrated through the inserts and settled on the tissue culture dish were fixed with 4% paraformaldehyde. Cell were then stained with hematoxylin and counted under a dissection microscope. Cells that migrated through the transwell inserts are compared for pCMV-neo and pCMV-1/3Nr-AS transfected cells.

Figure 19 shows the effect of anti-sense hNr-CAM overexpression on the invasion ability of 5GB cells. Briefly, $1x10^4$ pCMV-neo and pCMV-1/3Nr-AS transfected cells were plated on transwell inserts ($8\mu m$) coated with 825ng extracellular matrix (ECM) gel from Coastar (Cambridge, MA). Inserts were placed in 6 well tissue culture dishes

containing the appropriate growth media with serum. Cells that migrated through the inserts and settled on the tissue culture dish were fixed with 4% paraformaldehyde. Cell are then stained with hematoxylin and counted under a dissection microscope. Cells that migrated through the transwell inserts are compared for pCMV-neo and pCMV-1/3Nr-AS transfected cells.

Figures 20 (A-C) show the effect of UV radiation on antisense hNr-CAM transfected 5GB cells. Figures 20 (A and B) show pCMV-neo and pCMV-1/3Nr-AS transfected cells analyzed for apoptosis. Arrows indicate cells undergoing apoptosis. Figure 20C shows % of cells undergoing apoptosis after treatment with 100 units of UV radiation.

over-expression on the tumor forming ability of 5GB cells in vivo. Lower three mice were injected with pCMV-neo transfected 5GB cells. Top three mice were injected with pCMV-1/3Nr-CAM transfected 5GB cells. The tumors and the site of injection are indicated by arrows.

Figures 22 (A and B) show the effects of intra-20 tumoral inoculation of plasmid expressing antisense hNr-CAM. Figure 22 A illustrates results obtained using three athymic nude mice injected 72 days post-implantation with 5 GB cells with 50 μ g either pCMVneo (control animal (\blacksquare)) or pCMVneo 1/3 Nr-AS (two animals (\blacksquare and Δ). Arrows indicate days of

25 intratumoral injection. See Section 7.2.9. for details. Figure 22B shows results with four athymic nude mice implanted 3x3mm pieces of glioblastoma tumor. This tumor was generated previously by injecting lx10⁷ 5GB glioblastoma cells subcutaneously. 28 days post implantation, 50μg of either pCMVneo or pCMV1/3Nr-AS plasmids were mixed with DMRIE (liposomes) reagent (Gibco/BRL) and injected twice a week for four weeks around the tumor site. Tumor size was measured twice a week with a caliper and tumor volume was determined.

Arrow indicates the first day of intra-tumoral injection. See Section 7.2.9. for details.

Figures 23 (A-C) show the effect of antisense hNr-CAM on GB1690 glioblastoma cells. The full length clone for 5 hNr-CAM was provided by William J. Dreyer (California Institute of Technology). A PCR product of approximately 1360 bases (Nucleotide positions 1410-2746) was generated from the full length hNr-CAM clone using specific primers (5'TAGATACAACTAGTCAATGCCTCTAATGAATATGG ATA3' (SEQ. ID. No.: 6); and 5' AGATAGATCCGCGGAATAGTAAA TCCGATA GCCTTGTA3' (SEQ. The PCR product was then cloned into Spel and ID. No.: 7). SacII sites of pCMV-neo vector and was termed as pCMV-2/3Nr-The pCMV-neo or pCMV-2/3Nr-AS were then transfected into AS. GB1690 glioblastoma cell line and selected in G418. Figures 23 (A and B) show cell morphology of cells transfected with 15 pCMV-neo and pCMV-2/3Nr-AS, respectively. Arrows indicate spindle shape cells. Figure 23C shows a comparison of number of soft agar colonies formed by pCMV-neo and pCMV-2/3Nr-AS transfected GB1690 cells respectively.

Figure 24 shows the sequence identity analysis 20 between human and rat Nr-CAM nucleotide sequence.

Figures 25 (A and B) schematically show the packaging of infectious, replication-incompetent retroviral particles (Figure 25 A) and a retroviral vector (Figure 25B). The retroviral vector, containing the gene of interest (antisense <a href="https://hww.ncm.nc.ning.nc.ni

integrase), and env (coat glycoproteins). Virus released from this cell line contains the products of these genes (and is infectious), but lacks the genes themselves, thus

preventing retroviral production from subsequently infected cell lines. Figure 25B shows pLXSN retroviral vector that can be used for cloning antisense hNr-CAM gene.

Figures 26 (A and B) show the identification of differentially expressed genes in 5GB glioblastoma cells transfected with pCMVneo (Figure 26B) or pCMV1/3Nr-AS (Figure 26A) vectors. Differential hybridization was performed as described in Section 8.. White arrow in Figure 26A indicates novel cDNA (accession# H77485) and selectin gene is indicated by black arrow (See Figure 26B).

5. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the identification of a novel role of Nr-CAM in cell transformation and aberrant cellular proliferation. In particular, the present invention relates to the discovery of altered expression of Nr-CAM in a number of primary tumors and cell lines derived from tumors, in addition to, the altered expression of ligands for Nr-CAM. Further, the present invention relates, in part, to the Applicants' surprising discovery that the inhibition of Nr-20 CAM gene expression or the inhibition of Nr-CAM activity in transformed cells reverses the transformed phenotype.

The present invention encompasses compounds and methods for the detection of aberrant Nr-CAM gene expression or activity as a diagnostic or prognostic tool to indicate a transformed, pre-cancerous or cancerous cell phenotype. The present invention further encompasses compounds and methods for the detection of aberrant gene expression or activity as a diagnostic or prognostic tool to indicate a transformed, pre-cancerous or cancerous cell phenotype. The present invention also encompasses compounds and methods for the modulation of Nr-CAM gene expression or activity as a method of treating or preventing a transformed, pre-cancerous or cancerous cell phenotype. In this regard, the present

invention provides nucleotide sequences of Nr-CAM genes, and amino acid sequences of their encoded proteins. The invention further provides fragments and other derivatives, and analogs, of Nr-CAM proteins. Nucleic acids encoding such fragments or derivatives are also within the scope of the invention. The invention provides Nr-CAM nucleic acids and their encoded proteins of humans and related genes (homologs) in other species. In specific embodiments, the Nr-CAM nucleic acids and proteins are from vertebrates, or more particularly, mammals. In a preferred embodiment of the invention, the Nr-CAM nucleic acids and proteins are of human origin. Production of the foregoing nucleic acids, proteins and derivatives, e.g., by recombinant methods, is provided.

Nr-CAM is a gene identified by the method of the invention, that is expressed at high levels in glioblastoma multiforme tissue as well as certain others forms of tumors and cancers.

The invention also provides Nr-CAM derivatives and analogs of the invention which are functionally active, i.e., they are capable of displaying one or more functional

20 activities described herein associated with a full-length (wild-type) Nr-CAM protein. Such functional activities include, but are not limited to, antigenicity, i.e., ability to bind (or compete with Nr-CAM for binding) to an anti-Nr-CAM antibody, immunogenicity, i.e., ability to generate

25 antibody which binds to Nr-CAM, and ability to bind (or compete with Nr-CAM for binding) to a ligand for Nr-CAM. The invention further provides fragments (and derivatives and analogs thereof) of Nr-CAM which comprise one or more domains of the Nr-CAM protein. Antibodies to Nr-CAM, its derivatives and analogs, are additionally provided.

The present invention also provides therapeutic and diagnostic methods and compositions based on Nr-CAM proteins and nucleic acids and anti-Nr-CAM antibodies. The invention

30

provides for treatment of disorders of overproliferation (e.g., cancer and hyperproliferative disorders) by administering compounds that decrease Nr-CAM activity (e.g., antibodies, Nr-CAM antisense nucleic acids).

The invention also provides methods of treatment of disorders involving deficient cell proliferation or in which cell proliferation (growth) is otherwise desirable (e.g., growth deficiencies, degenerative disorders, lesions, physical trauma) by administering compounds that promote Nr-CAM function.

The present invention further provides screening assays to identify novel agents which target Nr-CAM gene or nucleic acid expression or Nr-CAM protein activity, including interaction with ligands, and, thus are potential therapeutic agents for the treatment or prevention of cell

- transformation, or pre-cancerous or cancerous phenotypes, i.e., tumorigenesis. The screening assays of the present invention may function to identify novel exogenous or endogenous agents that inhibit Nr-CAM expression or inhibit the interaction between Nr-CAM and its ligand. A variety of
- 20 protocols and techniques may be used to identify drugs that inhibit Nr-CAM expression and/or Nr-CAM activity, and as a result inhibit Nr-CAM participation in aberrant cellular proliferative activity. Such identified agents have utility in the treatment of hosts demonstrating a cellular transformed phenotype or aberrant cellular proliferative.
- transformed phenotype or aberrant cellular proliferative behavior, and advantageously would be effective to treat and/or prevent tumorigenesis.

The present invention further encompasses

pharmaceutical compositions containing the novel agents
identified by the screening assays described herein. The

invention provides therapeutic modalities and pharmaceutical
compositions for the treatment of tumorigenesis and the
prevention of transformed phenotypes. The therapeutic

modalities of the present invention further encompass combination therapies in which an agent which inhibits Nr-CAM expression and/or activity, and at least one other therapeutic agent, e.g., a chemotherapeutic agent, are administered either concurrently, e.g., as an admixture, separately but simultaneously or concurrently, or sequentially.

The novel therapeutic combinations of the present invention provide a means of treatment which may not only reduce the effective dose of either drug required for antitransformation or antitumorigenesis, thereby reducing toxicity, but may improve the absolute therapeutic effect as a result of attacking aberrant cellular proliferation through a variety of mechanisms.

The invention is illustrated by way of examples

15 infra which disclose, inter alia, the isolation and
characterization of Nr-CAM, and patterns of expression of NrCAM in certain tumors (see Section 6).

For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the subsections which follow.

5.1. IDENTIFICATION OF ROLE OF Nr-CAM IN TRANSFORMATION

The present invention relates to a novel role of Nr-CAM in the promotion of cell transformation and tumorigenesis. In particular, the present invention relates to the Applicants' findings that (a) Nr-CAM is highly over-expressed in glioblastoma multiforme tumor tissue and is over-expressed a number of other primary tumors; and (b) over-expression of Nr-CAM in the anti-sense orientation results in decreased cellular proliferation and colony formation of glioblastoma cells in soft agar.

The present invention further relates to the Applicants' findings that Nr-CAM is over-expressed in several

brain tumor derived cell lines and primary brain tumor tissues, including astrocytoma grade IV, glioma, glioblastoma and neuroectodermal human tumor cell lines compared to NBT. Low or no expression of hNr-CAM was observed in cell lines derived from astrocytoma III, medulloblastoma, neuroblastoma and NBT. Further, Nr-CAM was found to be expressed at high levels in melanoma G361, lymphoblastic leukemia cell lines, and Burkitt's lymphoma cell lines. A low level of hNr-CAM expression was observed in colorectal adenocarcinoma, lung carcinoma, pro-myelocytic leukemia HeLa cell S3, and chronic myelogenous leukemia.

The present invention relates to the role of Nr-CAM in promotion of cell transformation and tumorigenesis, and provides methods including the use of Nr-CAM nucleic acids and nucleic acids which hybridize or complement Nr-CAM nucleic acids, as diagnostic and prognostic tools for the detection of transformed, pre-cancerous and cancerous phenotypes. The present invention provides methods for use of Nr-CAM nucleic acids and those which complement and/or hybridize to nucleic acid sequences which encode Nr-CAM as therapeutics to treat, inhibit or prevent transformed, pre-cancerous and cancerous phenotypes. In particular, the invention provides compositions comprising nucleic acid sequences which inhibit Nr-CAM expression as therapeutics to treat or prevent transformed, pre-cancerous, and cancerous phenotypes.

5.2. THE PRODUCTION OF Nr-CAM NUCLEIC ACIDS, POLYPEPTIDES AND ANTIBODIES AS DIAGNOSTICS, THERAPEUTICS AND COMPONENTS FOR SCREENING ASSAYS

The present invention encompasses the use of agents for the detection of aberrant Nr-CAM gene expression as diagnostic or prognostic tools to detect a transformed phenotype, pre-cancerous or cancerous condition. Diagnostic

or prognostic tools which may be used in accordance with the present invention include, but are not limited to, (a) nucleic acids which hybridize or are complementary to the Nr-CAM nucleotide sequence; (b) polypeptides, peptide fragments or synthetic molecules which bind to the Nr-CAM ligand binding domain; and (c) antibodies which bind to Nr-CAM.

which inhibit Nr-CAM expression and/or protein activity as therapeutics for the treatment and/or prevention of a transformed or pre-cancerous phenotype, or cancer or tumorigenesis. Therapeutic agents which may be used in accordance with the present invention include, but are not limited to, (a) nucleic acids which inhibit Nr-CAM gene expression, e.g., antisense molecules, ribozymes or triple helix molecules complementary to Nr-CAM; (b) polypeptides, peptides, antibodies, small organic molecules or synthetic molecules which inhibit Nr-CAM activity or prevent Nr-CAM from binding its ligand; and (c) peptides, polypeptides, antibodies, small organic molecules or synthetic molecules which act as antagonists of Nr-CAM activity.

The present invention provides screening assays for the identification of agents which inhibit Nr-CAM gene expression and/or activity. In one embodiment of the invention, an important component of the screening assays of the present invention are nucleotide coding sequences encoding Nr-CAM proteins, polypeptides and peptides. The present invention further encompasses (a) DNA vectors that contain any of the foregoing Nr-CAM encoding sequences and/or their complements; (b) DNA expression vectors that contain any of the foregoing Nr-CAM coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (c) genetically engineered host cells that contain any of the foregoing Nr-CAM coding sequences operatively associated with

a regulatory element that directs the expression of the coding sequences in the host cell.

5.2.1. THE Nr-CAM NUCLEIC ACIDS

The invention relates to the nucleotide sequences 5 of Nr-CAM nucleic acids. In an embodiment, the Nr-CAM nucleic acids comprise the nucleotide sequence shown in Figure 2A, i.e., SEQ. ID. No.: 1 or specific regions thereof. In specific embodiments, Nr-CAM nucleic acids comprise the cDNA sequences of SEQ. ID. NO.: 5, or the coding regions thereof, or nucleotide sequences acids encoding a Nr-CAM protein (e.g., a protein having the sequence shown in Figure 2B, i.e., SEQ. ID. NO.: 2) or a fragment thereof. Nucleic acids of the present invention can be single or double stranded. The invention also relates to nucleic acids hybridizable to or complementary to the foregoing sequences. In specific aspects, nucleic acids are provided which comprise a sequence complementary to at least 10, 25, 50, 100, 200, or 250 contiguous nucleotides of a Nr-CAM gene. In a specific embodiment, a nucleic acid which is hybridizable 20 to a Nr-CAM nucleic acid (e.g., having sequence SEQ. ID. NO.: 1), or to a nucleic acid encoding a Nr-CAM derivative, under conditions of low stringency is provided.

By way of example and not limitation, procedures using such conditions of low stringency are as follows (see also Shilo and Weinberg, 1981, Proc. Natl. Acad. Sci. USA 78:6789-6792): Filters containing DNA are pretreated for 6 h at 40°C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 μg/ml denatured salmon sperm DNA. Hybridizations are carried out in the same solution with the following modifications: 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 μg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and 5-20 X 106 cpm ³²P-labeled probe is used. Filters are

incubated in hybridization mixture for 18-20 h at 40°C, and then washed for 1.5 h at 55°C in a solution containing 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash solution is replaced with fresh solution and incubated an additional 1.5 h at 60°C. Filters are blotted dry and exposed for autoradiography. If necessary, filters are washed for a third time at 65-68°C and re-exposed to film. Other conditions of low stringency which may be used are well known in the art (e.g., as employed for cross-species hybridizations).

In another specific embodiment, -a-nucleic acid which is hybridizable to a Nr-CAM nucleic acid under conditions of high stringency is provided. By way of example and not limitation, procedures using such conditions of high stringency are as follows: Prehybridization of filters 15 containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 μ g/ml denatured salmon sperm DNA. Filters are hybridized for 48 h at 65°C in prehybridization mixture containing 100 µg/ml 20 denatured salmon sperm DNA and 5-20 X 106 cpm of 32P-labeled probe. Washing of filters is done at 37°C for 1 h in a solution containing 2X SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA. This is followed by a wash in 0.1X SSC at 50°C for 45 min before autoradiography. Other conditions of high stringency which may be used are well known in the art.

In another specific embodiment, a nucleic acid which is hybridizable to a Nr-CAM nucleic acid under conditions of moderate stringency is provided.

25

Various other stringency conditions which promote nucleic acid hybridization can be used. For example,

30 hybridization in 6x SSC at about 45° C, followed by washing in 2xSSC at 50° C may be used. Alternatively, the salt concentration in the wash step can range from low stringency

of about 5xSSC at 50°C, to moderate stringency of about 2xSSC at 50°C, to high stringency of about 0.2x SSC at 50°C. In addition, the temperature of the wash step can be increased from low stringency conditions at room temperature, to moderately stringent conditions at about 42°C, to high stringency conditions at about 65°C. Other conditions include, but are not limited to, hybridizing at 68°C in 0.5M NaHPO4 (pH7.2) / 1 mM EDTA / 7% SDS, or hybridization in 50% formamide / 0.25M NaHPO4 (pH 7.2) / .25 M NaCl / 1 mM EDTA / 7% SDS; followed by washing in 40 mM NaHPO4 (pH 7.2) / 1 mM EDTA / 5% SDS at 42°C or in 40 mM NaHPO4 (pH7.2) 1 mM EDTA / 1% SDS at 50°C. Both temperature and salt may be varied, or alternatively, one or the other variable may remain constant while the other is changed.

Low, moderate and high stringency conditions are

15 well known to those of skill in the art, and will vary
predictably depending on the base composition of the
particular nucleic acid sequence and on the specific organism
from which the nucleic acid sequence is derived. For
guidance regarding such conditions see, for example, Sambrook
20 et al., 1989, Molecular Cloning, A Laboratory Manual, Second
Edition, Cold Spring Harbor Press, N.Y., pp. 9.47-9.57; and
Ausubel et al., 1989, Current Protocols in Molecular Biology,
Green Publishing Associates and Wiley Interscience, N.Y.

The invention also encompasses nucleic acids having at least 60%, 70%, 75%, 80%, 90%, 95% or greater sequence identity when compared to a portion of identical-sized hNr-CAM sequence shown in Figure 2A or when compared to said sequence when the alignment or comparison is conducted by a computer homology programmer search aligorithm known in the art.

By way of example and not limitation, useful computer homology programs include the following: Basic Local Alignment Search Tool (BLAST) (www.ncbi.nlm.nih.gov)

(Altschul et al., 1990, J. of Molec. Biol., 215:403-410, "The BLAST Algorithm"; FASTA (also TFASTA) (see, Pearson et al., 1988, Proc. Nat'l Acad. Sci. USA 85:2444-2448) and CLUSTALW (see, Higgins et al., 1996, Methods Enzymol 266:383-402).

Altschul et al., 1997, Nuc. Acids Res.
25:3389-3402) describe BLAST, a heuristic search algorithm
tailored to searching for sequence similarity which ascribes
significance using the statistical methods of Karlin and
Altschul 1990, Proc. Nat'l Acad. Sci. USA, 87:2264-68; 1993,

Proc. Nat'l Acad. Sci. USA 90:5873-77. Five specific BLAST
programs perform the following tasks:

- 1) The BLASTP program compares an amino acid query sequence against a protein sequence database.
- 2) The BLASTN program compares a nucleotide query sequence against a nucleotide sequence database.
- 3) The BLASTX program compares the six-frame conceptual translation products of a nucleotide query sequence (both strands) against a protein sequence database.
- 4) The TBLASTN program compares a protein query sequence against a nucleotide sequence database translated in 20 all six reading frames (both strands).
 - 5) The TBLASTX program compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

As will be understood by those skilled in the art,
the TBLASTN program is particularly useful to identify
nucleic acids with a desired percent identity and the BLASTP
program is particularly useful to identify amino acid
sequences with a desired percent identity.

Smith-Waterman (database: European Bioinformatics Institute wwwz.ebi.ac.uk/bic_sw/) (Smith-Waterman, 1981, J. 30 of Molec. Biol., 147:195-197) is a mathematically rigorous algorithm for sequence alignments.

FASTA (see Pearson et al., 1988, Proc. Nat'l Acad. Sci. USA, 85:2444-2448) is a heuristic approximation to the Smith-Waterman algorithm. For a general discussion of the procedure and benefits of the BLAST, Smith-Waterman and FASTA algorithms see Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" (www.psc.edu) and references cited therein.

Nucleic acids encoding derivatives and analogs of Nr-CAM proteins (see Sections 5.2.2), and Nr-CAM antisense nucleic acids are additionally provided. As is readily apparent, as used herein, a "nucleic acid encoding a fragment or portion of a Nr-CAM protein" shall be construed as referring to a nucleic acid encoding only the recited fragment or portion of the Nr-CAM protein and not the other contiguous portions of the Nr-CAM protein as a continuous sequence.

Fragments of Nr-CAM nucleic acids comprising regions conserved between other Nr-CAM nucleic acids, of the same or different species, are also provided. Nucleic acids encoding one or more Nr-CAM domains are provided.

Specific embodiments for the cloning of a Nr-CAM gene, presented as a particular example but not by way of limitation, follow:

For expression cloning (a technique commonly known in the art), an expression library is constructed by methods known in the art. For example, mRNA (e.g., human) is isolated, cDNA is made and ligated into an expression vector (e.g., a bacteriophage derivative) such that it is capable of being expressed by the host cell into which it is then introduced. Various screening assays can then be used to select for the expressed Nr-CAM product. In one embodiment, anti-Nr-CAM antibodies can be used for selection.

In another embodiment, polymerase chain reaction (PCR) is used to amplify the desired sequence in a genomic or

cDNA library, prior to selection. Oligonucleotide primers representing known Nr-CAM sequences can be used as primers in In a preferred aspect, the oligonucleotide primers represent at least part of the Nr-CAM sequence presented in 5 Figure 2A. The synthetic oligonucleotides may be utilized as primers to amplify by PCR sequences from a source (RNA or DNA), preferably a cDNA library, of potential interest. PCR can be carried out, e.g., by use of a Perkin-Elmer Cetus thermal cycler and Taq polymerase (Gene Amp™). The DNA being amplified can include mRNA, cDNA, or genomic DNA from any 10 eukaryotic species. One can choose to synthesize several different degenerate primers, for use in the PCR reactions. It is also possible to vary the stringency of hybridization conditions used in priming the PCR reactions, to allow for greater or lesser degrees of nucleotide sequence similarity 15 between the known Nr-CAM nucleotide sequence and the nucleic acid homolog being isolated. For cross species hybridization, low stringency conditions are preferred. same species hybridization, moderately stringent conditions are preferred. After successful amplification of a segment 20 of a Nr-CAM homolog, that segment may be molecularly cloned and sequenced, and utilized as a probe to isolate a complete cDNA or genomic clone. This, in turn, will permit the determination of the gene's complete nucleotide sequence, the analysis of its expression, and the production of its protein product for functional analysis, as described infra. fashion, additional genes encoding Nr-CAM proteins and Nr-CAM analogs may be identified.

The above-methods are not meant to limit the following general description of methods by which clones of Nr-CAM may be obtained.

Any eukaryotic cell potentially can serve as the nucleic acid source for the molecular cloning of the Nr-CAM gene. The nucleic acid sequences encoding Nr-CAM can be

30

isolated from vertebrate sources, including mammalian sources, such as porcine, bovine, feline, and equine, canine, human, as well as additional primate sources, avian, reptilian, amphibian, piscine, etc. sources, non-vertebrate 5 sources such as insects, from plants, etc. The DNA may be obtained by standard procedures known in the art from cloned DNA (e.g., a DNA "library"), by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell. (See, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; Glover, D.M. (ed.), 1985, DNA Cloning: A Practical Approach, MRL Press, Ltd., Oxford, U.K. Vol. I, II.) Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding 15 regions; clones derived from cDNA will contain only exon sequences. Whatever the source, the gene should be molecularly cloned into a suitable vector for propagation of the gene.

In the molecular cloning of the gene from genomic DNA, DNA fragments are generated, some of which will encode the desired gene. The DNA may be cleaved at specific sites using various restriction enzymes. Alternatively, one may use DNAse in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, as for example, by sonication. The linear DNA fragments can then be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing the desired gene may be accomplished in a number of ways. For example, if an amount of a portion of a Nr-CAM (of any species) gene or its specific RNA, or a fragment thereof (see

Section 5.6), is available and can be purified and labeled, the generated DNA fragments may be screened by nucleic acid hybridization to the labeled probe (Benton and Davis, 1977, Science 196:180; Grunstein and Hogness, 1975, Proc. Natl.

- Acad. Sci. U.S.A. 72:3961). Those DNA fragments with substantial homology to the probe will hybridize. It is also possible to identify the appropriate fragment by restriction enzyme digestion(s) and comparison of fragment sizes with those expected according to a known restriction map if such is available. Further selection can be carried out on the basis of the properties of the gene. Alternatively, the presence of the gene may be detected by assays based on the physical, chemical, or immunological properties of its
- which hybrid-select the proper mRNAs, can be selected which produce a protein that, e.g., has similar or identical electrophoretic migration, isoelectric focusing behavior, proteolytic digestion maps, promotion of cell proliferation activity, substrate binding activity, or antigenic properties of Nr-CAM. If an antibody to Nr-CAM is available, the Nr-CAM

expressed product. For example, cDNA clones, or DNA clones

20 protein may be identified by binding of labeled antibody to the putatively Nr-CAM synthesizing clones, in an ELISA (enzyme-linked immunosorbent assay)-type procedure.

The Nr-CAM gene can also be identified by mRNA selection by nucleic acid hybridization followed by in vitro translation. In this procedure, fragments are used to isolate complementary mRNAs by hybridization. Such DNA fragments may represent available, purified Nr-CAM DNA of another species (e.g., human, mouse, etc.).

Immunoprecipitation analysis or functional assays (e.g.,

aggregation ability in vitro; binding to receptor; see infra) of the in vitro translation products of the isolated products of the isolated mRNAs identifies the mRNA and, therefore, the complementary DNA fragments that contain the desired

sequences. In addition, specific mRNAs may be selected by adsorption of polysomes isolated from cells to immobilized antibodies specifically directed against Nr-CAM protein. A radiolabelled Nr-CAM cDNA can be synthesized using the selected mRNA (from the adsorbed polysomes) as a template. The radiolabelled mRNA or cDNA may then be used as a probe to identify the Nr-CAM DNA fragments from among other genomic DNA fragments.

Alternatives to isolating the Nr-CAM genomic DNA include, but are not limited to, chemically synthesizing the gene sequence itself from a known sequence or making cDNA to the mRNA which encodes the Nr-CAM protein. For example, RNA for cDNA cloning of the Nr-CAM gene can be isolated from cells which express Nr-CAM. Other methods are possible and within the scope of the invention.

15 The identified and isolated gene can then be inserted into an appropriate cloning vector. A large number of vector-host systems known in the art may be used. Possible vectors include, but are not limited to, plasmids or modified viruses, but the vector system must be compatible 20 with the host cell used. Such vectors include, but are not limited to, bacteriophages such as lambda derivatives, or plasmids such as PBR322 or pUC plasmid derivatives or the Bluescript vector (Stratagene). The insertion into a cloning vector can, for example, be accomplished by ligating the DNA fragment into a cloning vector which has complementary 25 cohesive termini. However, if the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules may be enzymatically modified. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; 30 these ligated linkers may comprise specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. In an alternative

method, the cleaved vector and Nr-CAM gene may be modified by homopolymeric tailing. Recombinant molecules can be introduced into host cells via transformation, transfection, infection, electroporation, etc., so that many copies of the gene sequence are generated.

In an alternative method, the desired gene may be identified and isolated after insertion into a suitable cloning vector in a "shot gun" approach. Enrichment for the desired gene, for example, by size fractionization, can be done before insertion into the cloning vector.

10

30

In specific embodiments, transformation of host cells with recombinant DNA molecules that incorporate the isolated Nr-CAM gene, cDNA, or synthesized DNA sequence enables generation of multiple copies of the gene. Thus, the gene may be obtained in large quantities by growing

15 transformants, isolating the recombinant DNA molecules from the transformants and, when necessary, retrieving the inserted gene from the isolated recombinant DNA.

The Nr-CAM sequences provided by the present invention include those nucleotide sequences encoding substantially the same amino acid sequences as found in native Nr-CAM proteins, and those encoded amino acid sequences with functionally equivalent amino acids, as well as those encoding other Nr-CAM derivatives or analogs, as described in Section 5.2.2 infra for Nr-CAM derivatives and analogs.

The Nr-CAM sequences provided by the present invention include those that encode Nr-CAM mutants that are constitutively expressed.

5.2.2. EXPRESSION OF Nr-CAM NUCLEIC ACIDS

The nucleotide sequence coding for a Nr-CAM protein or a functionally active analog or fragment or other derivative thereof, can be inserted into an appropriate

expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. The necessary transcriptional and translational signals can also be 5 supplied by the native Nr-CAM gene and/or its flanking regions. A variety of host-vector systems may be utilized to express the protein-coding sequence. These include but are not limited to mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g., baculovirus); microorganisms such 10 as yeast containing yeast vectors, or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation 15 elements may be used. In specific embodiments, the human Nr-CAM gene is expressed, or a sequence encoding a functionally active portion of human Nr-CAM. In yet another embodiment, a fragment of Nr-CAM comprising a domain of the Nr-CAM protein is expressed.

Any of the methods previously described for the insertion of DNA fragments into a vector may be used to construct expression vectors containing a chimeric gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include in vitro recombinant DNA and synthetic techniques and in vivo recombinants (genetic recombination). Expression of nucleic acid sequence encoding a Nr-CAM protein or peptide fragment may be regulated by a second nucleic acid sequence so that the Nr-CAM protein or peptide is expressed in a host transformed with the recombinant DNA molecule. For example, expression of a Nr-CAM protein may be controlled by any promoter/enhancer element known in the art. Promoters which may be used to control Nr-CAM expression include, but

are not limited to, the SV40 early promoter region (Bernoist and Chambon, 1981, Nature 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, Cell 22:787-797), the herpes 5 thymidine kinase promoter (Wagner et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, Nature 296:39-42); prokaryotic expression vectors such as the β lactamase promoter (Villa-Kamaroff, et al., 1978, Proc. Natl. Acad. Sci. U.S.A. <u>75</u>:3727-3731), or the tac promoter (DeBoer, et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:21-25); see also "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242:74-94; plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella et al., 1983, Nature 303:209-213) or the 15 cauliflower mosaic virus 35S RNA promoter (Gardner, et al., 1981, Nucl. Acids Res. 9:2871), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., 1984, Nature 310:115-120); promoter elements from yeast or other fungi such as the Gal 4 20 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control 25 region which is active in pancreatic acinar cells (Swift et al., 1984, Cell 38:639-646; Ornitz et al., 1986, Cold Spring Harbor Symp. Quant. Biol. 50:399-409; MacDonald, 1987, Hepatology 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, Nature 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, Cell 38:647-658; Adames et al., 1985, Nature 318:533-538; Alexander et al., 1987, Mol. Cell. Biol. 7:1436-1444), mouse

mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, Cell 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, Genes and Devel.

- 5 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, Mol. Cell. Biol. 5:1639-1648; Hammer et al., 1987, Science 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., 1987, Genes and Devel. 1:161-171), beta-
- globin gene control region which is active in myeloid cells

 (Mogram et al., 1985, Nature 315:338-340; Kollias et al.,
 1986, Cell 46:89-94; myelin basic protein gene control region
 which is active in oligodendrocyte cells in the brain
 (Readhead et al., 1987, Cell 48:703-712); myosin light chain2 gene control region which is active in skeletal muscle
- (Sani, 1985, Nature 314:283-286), and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, Science 234:1372-1378).

In a specific embodiment, a vector is used that comprises a promoter operably linked to a Nr-CAM-encoding 20 nucleic acid, one or more origins of replication, and, optionally, one or more selectable markers (e.g., an antibiotic resistance gene).

In a specific embodiment, an expression construct is made by subcloning a Nr-CAM coding sequence into the EcoRI restriction site of each of the three pGEX vectors (Glutathione S-Transferase expression vectors; Smith and Johnson, 1988, Gene 7:31-40). This allows for the expression of the Nr-CAM protein product from the subclone in the correct reading frame.

In another specific embodiment, the promoter that is operably linked to the hNr-CAM gene is not the native hNr-CAM gene promoter (i.e., it is a heterologous promoter).

Expression vectors containing Nr-CAM gene inserts can be identified by three general approaches: (a) nucleic acid hybridization, (b) presence or absence of "marker" gene functions, and (c) expression of inserted sequences. In the 5 first approach, the presence of a Nr-CAM gene inserted in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted Nr-CAM gene. In the second approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (e.g., thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of a Nr-CAM gene in the vector. For example, if the Nr-CAM gene is inserted within the marker 15 gene sequence of the vector, recombinants containing the Nr-CAM insert can be identified by the absence of the marker gene function. In the third approach, recombinant expression vectors can be identified by assaying the Nr-CAM product expressed by the recombinant. Such assays can be based, for 20 example, on the physical or functional properties of the Nr-CAM protein in in vitro assay systems, e.g., binding with anti-Nr-CAM antibody, promotion of cell proliferation.

Once a particular recombinant DNA molecule is identified and isolated, several methods known in the art may be used to propagate it. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As previously explained, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as

baculovirus; yeast vectors; bacteriophage vectors (e.g.,

lambda), and plasmid and cosmid DNA vectors, to name but a few.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or ς modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus, expression of the genetically engineered Nr-CAM protein may be controlled. Furthermore, different host cells have characteristic and specific mechanisms for the translational 10 and post-translational processing and modification (e.g., glycosylation, phosphorylation of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system can be used to 15 produce an unglycosylated core protein product. Expression in yeast will produce a glycosylated product. Expression in mammalian cells can be used to ensure "native" glycosylation of a heterologous protein. Furthermore, different vector/host expression systems may effect processing 20 reactions to different extents.

In other specific embodiments, the Nr-CAM protein, fragment, analog, or derivative may be expressed as a fusion, or chimeric protein product (comprising the protein, fragment, analog, or derivative joined via a peptide bond to a heterologous protein sequence (of a different protein)).

Such a chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other by methods known in the art, in the proper coding frame, and expressing the chimeric product by methods commonly known in the art. Alternatively, such a chimeric product may be made by protein synthetic techniques, e.g., by use of a peptide synthesizer.

Both cDNA and genomic sequences can be cloned and expressed.

5.2.3. IDENTIFICATION AND PURIFICATION OF THE Nr-CAM PRODUCTS

In particular aspects, the invention provides amino acid sequences of Nr-CAM, preferably human Nr-CAM, and fragments and derivatives thereof which comprise an antigenic determinant (i.e., can be recognized by an antibody) or which are otherwise functionally active, as well as nucleic acid sequences encoding the foregoing. "Functionally active" Nr-CAM material as used herein refers to that material displaying one or more functional activities associated with a full-length (wild-type) Nr-CAM protein, e.g., promotion of cell proliferation, binding to a Nr-CAM substrate or Nr-CAM binding partner, antigenicity (binding to an anti-Nr-CAM antibody), immunogenicity, etc.

In other specific embodiments, the invention provides fragments of a Nr-CAM protein consisting of at least 6 amino acids, 10 amino acids, 50 amino acids, or of at least 75 amino acids of SEQ. ID. No.: 2. In other embodiments, the invention provides proteins comprising, having, or consisting essentially of a sequence of amino acids 100% identical with SEQ. ID. NO.: 2, SEQ. ID. NO.: 2 or a protein encoded by SEQ. ID. NO.: 1, or any combination of the foregoing. Fragments or proteins comprising such sequences are particularly advantageously used for immunotherapy as described below. Fragments, or proteins comprising fragments, lacking some or all of the foregoing regions of a Nr-CAM protein are also provided. Nucleic acids encoding the foregoing are provided. In specific embodiments, the foregoing proteins or fragments are not more than 25, 50 or 100 contiguous amino acids.

Once a recombinant which expresses the Nr-CAM gene sequence is identified, the gene product can be analyzed.

This is achieved by assays based on the physical or functional properties of the product, including radioactive labelling of the product followed by analysis by gel electrophoresis, immunoassay, etc.

Once the Nr-CAM protein is identified, it may be isolated and/or purified by standard methods including chromatography (e.g., ion exchange, affinity, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. The functional properties may be evaluated using any suitable assay (see Section 5.3).

Alternatively, once a Nr-CAM protein produced by a recombinant is identified, the amino acid sequence of the protein can be deduced from the nucleotide sequence of the chimeric gene contained in the recombinant. As a result, the protein can be synthesized by standard chemical methods known in the art (e.g., see Hunkapiller, M., et al., 1984, Nature 310:105-111).

In another alternate embodiment, native Nr-CAM proteins can be purified from natural sources, by standard 20 methods such as those described above (e.g., immunoaffinity purification).

In a specific embodiment of the present invention, such Nr-CAM proteins, whether produced by recombinant DNA techniques or by chemical synthetic methods or by purification of native proteins, include but are not limited to those containing, as a primary amino acid sequence, all or part of the amino acid sequence substantially, as well as fragments and other derivatives, and analogs as shown in Figure 2B (SEQ. ID. NO.: 2) thereof, including proteins homologous thereto.

5.2.4. ANTIBODIES AND IMMUNE CELLS TO Nr-CAM

5.2.4.1. GENERATION OF ANTIBODIES TO Nr-CAM PROTEINS AND DERIVATIVES THEREOF

fragments or other derivatives, or analogs thereof, may be used as an immunogen to generate antibodies which immunospecifically bind such an immunogen. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression library. In a specific embodiment, antibodies to a human Nr-CAM protein are produced. In another embodiment, antibodies to a domain of a Nr-CAM protein are produced. In a specific embodiment, fragments of a Nr-CAM protein identified as hydrophilic are used as immunogens for antibody production.

In another specific embodiment, the antibody to a

15 human Nr-CAM protein is a bispecific antibody (see generally,
e.g. Fanger and Drakeman, 1995, Drug News and Perspectives 8:
133-137). Such a bispecific antibody is genetically
engineered to recognize both (1) a human Nr-CAM epitope and
(2) one of a variety of "trigger" molecules, e.g. Fc

receptors on myeloid cells, and CD3 and CD2 on T cells, that
have been identified as being able to cause a cytotoxic Tcell to destroy a particular target. Such bispecific
antibodies can be prepared either by chemical conjugation,
hybridoma, or recombinant molecular biology techniques known
to the skilled artisan.

Various procedures known in the art may be used for the production of polyclonal antibodies to a Nr-CAM protein or derivative or analog. In a particular embodiment, rabbit polyclonal antibodies to an epitope of a Nr-CAM protein, or a subsequence thereof, can be obtained. For the production of antibody, various host animals can be immunized by injection with the native Nr-CAM protein, or a synthetic version, or derivative (e.g., fragment) thereof, including but not

limited to rabbits, mice, rats, etc. Various adjuvants may be used to increase the immunological response, depending on the host species, and including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum.

For preparation of monoclonal antibodies directed 10 toward a Nr-CAM protein sequence or analog thereof, any technique which provides for the production of antibody molecules by continuous cell lines in culture may be used. For example, the hybridoma technique originally developed by Kohler and Milstein (1975, Nature 256:495-497), as well as 15 the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, Immunology Today $\underline{4}$:72), and the EBVhybridoma technique to produce human monoclonal antibodies (Cole et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). In an additional 20 embodiment of the invention, monoclonal antibodies can be produced in germ-free animals utilizing technology described in PCT/US90/02545. According to the invention, human antibodies may be used and can be obtained by using human hybridomas (Cote et al., 1983, Proc. Natl. Acad. Sci. U.S.A. 80:2026-2030) or by transforming human B cells with EBV virus in vitro (Cole et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, pp. 77-96). In fact, according to the invention, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, PROC. NATL. ACAD. SCI. U.S.A. <u>81</u>:6851-6855; Neuberger et al., 1984, Nature 312:604-608; Takeda et al., 1985, Nature 314:452-454) by splicing the genes from a mouse antibody molecule specific for Nr-CAM together with genes from a human antibody molecule

of appropriate biological activity can be used; such antibodies are within the scope of this invention.

According to the invention, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce Nr-CAM-specific single chain antibodies. An additional embodiment of the invention utilizes the techniques described for the construction of Fab expression libraries (Huse et al., 1989, Science 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for Nr-CAM proteins, derivatives, or analogs.

Antibody fragments which contain the idiotype of the molecule can be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, the Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent, and Fv fragments.

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, e.g. ELISA (enzyme-linked immunosorbent assay). For example, to select antibodies which recognize a specific domain of a Nr-CAM protein, one may assay generated hybridomas for a product which binds to a Nr-CAM fragment containing such domain. For selection of an antibody that specifically binds a first Nr-CAM homolog but which does not specifically bind a different Nr-CAM homolog, one can select on the basis of positive binding to the first Nr-CAM homolog and a lack of binding to the second Nr-CAM homolog.

Antibodies specific to a domain of a Nr-CAM protein are also provided.

30

The foregoing antibodies can be used in methods known in the art relating to the localization and activity of the Nr-CAM protein sequences of the invention, e.g., for imaging these proteins, measuring levels thereof in ς appropriate physiological samples, in diagnostic methods, etc.

In another embodiment of the invention (see infra), anti-Nr-CAM antibodies and fragments thereof containing the binding domain are Therapeutics.

10 Antibodies and antigen-binding antibody fragments may also be conjugated to a heterologous protein or peptide by chemical conjugation or recombinant DNA technology. resultant chimeric protein possesses the antigen-binding specificity of the antibody and the function of the 15 heterologous protein. For example, a polynucleotide encoding the antigen binding region of an antibody specific for the extracellular domain of Nr-CAM can be genetically fused to a coding sequence for the zeta chain of the T cell receptor. After expressing this construct in T cells, the T cells are 20 expanded ex vivo and infused into a brain cancer patient. cells expressing this chimeric protein are specifically directed to tumors that express Nr-CAM as a result of the antibody binding specificity and cause tumor cell killing. Alternatively, an antibody is fused to a protein which induces migration of leukocytes or has an affinity to attract 25 other compounds to a tumor cite. A specific protein of this type is streptavidin. The binding of a streptavidin conjugated antibody to a tumor cell can be followed by the addition of a biotinylated drug, toxin or radioisotope to cause tumor specific killing.

30 Kits for use with such in vitro tumor localization and therapy methods containing the monoclonal antibodies (or fragments thereof) conjugated to any of the above types of

substances can be prepared. The components of the kits can be packaged either in aqueous medium or in lyophilized form. When the monoclonal antibodies (or fragments thereof) are used in the kits in the form of conjugates in which a label or a therapeutic moiety is attached, such as a radioactive metal ion or a therapeutic drug moiety, the components of such conjugates can be supplied either in fully conjugated form, in the form of intermediates or as separate moieties to be conjugated by the user of the kit.

10 5.2.5. Nr-CAM PROTEINS, DERIVATIVES AND ANALOGS

The invention further encompasses compositions comprising Nr-CAM proteins, and derivatives (including but not limited to fragments) and analogs of Nr-CAM proteins, in particular, those derivatives which act as antagonists of Nr-15 CAM activity. Nucleic acids encoding Nr-CAM protein derivatives and protein analogs are also provided. In one embodiment, the Nr-CAM proteins are encoded by the Nr-CAM nucleic acids described in Section 5.2.1. supra. In particular aspects, the proteins, derivatives, or analogs are of Nr-CAM proteins of animals, e.g., fly, frog, mouse, rat, pig, cow, dog, monkey, human, or of plants.

related to Nr-CAM are within the scope of the present invention. In a specific embodiment, the derivative or analog is functionally active, i.e., capable of exhibiting one or more functional activities associated with a full-length, wild-type Nr-CAM protein. As one example, such derivatives or analogs which have the desired immunogenicity or antigenicity can be used, for example, in immunoassays, for immunization, for inhibition of Nr-CAM activity, etc.

30 Derivatives or analogs that retain, or alternatively lack or

inhibit, a desired Nr-CAM property of interest (e.g., binding to Nr-CAM binding partner, promotion of cell proliferation),

can be used as inducers, or inhibitors, respectively, of such property and its physiological correlates. A specific embodiment relates to a Nr-CAM fragment that can be bound by an anti-Nr-CAM antibody. Derivatives or analogs of Nr-CAM 5 can be tested for the desired activity by procedures known in the art, including but not limited to the assays described in Sections 5.3 and 5.5.

In particular, Nr-CAM derivatives can be made by altering Nr-CAM sequences by substitutions, additions or deletions that provide for functionally equivalent molecules. Due to the degeneracy of nucleotide coding sequences, other DNA sequences which encode substantially the same amino acid sequence as a Nr-CAM gene may be used in the practice of the present invention. These include but are not limited to nucleotide sequences comprising all or portions of Nr-CAM 15 genes which are altered by the substitution of different codons that encode a functionally equivalent amino acid residue within the sequence, thus producing a silent change. Likewise, the Nr-CAM derivatives of the invention include, but are not limited to, those containing, as a primary amino 20 acid sequence, all or part of the amino acid sequence of a Nr-CAM protein including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a silent For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, 30 leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine,

25

asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

- In a specific embodiment of the invention, proteins consisting of or comprising a fragment of a Nr-CAM protein consisting of at least 10 (continuous) amino acids of the Nr-CAM protein is provided. In other embodiments, the fragment consists of at least 20 or 50 amino acids of the Nr-CAM protein. In specific embodiments, such fragments are not larger than 35, 100 or 200 amino acids. Derivatives or
- larger than 35, 100 or 200 amino acids. Derivatives or analogs of Nr-CAM include but are not limited to those molecules comprising regions that are substantially homologous to Nr-CAM or fragments thereof (e.g., in various embodiments, at least 60% or 70% or 80% or 90% or 95%
- identity over an amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a computer homology program known in the art) or whose encoding nucleic acid is capable of hybridizing to a coding Nr-CAM sequence, under stringent, moderately
- 20 stringent, or nonstringent conditions. See, supra Section
 5.2.1. for useful computer programs for sequence comparisons.

The Nr-CAM derivatives and analogs of the invention can be produced by various methods known in the art. The manipulations which result in their production can occur at the gene or protein level. For example, the cloned Nr-CAM gene sequence can be modified by any of numerous strategies known in the art (Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York). The sequence can be cleaved at appropriate sites with restriction

endonuclease(s), followed by further enzymatic modification if desired, isolated, and ligated in vitro. In the production of the gene encoding a derivative or analog of Nr-

CAM, care should be taken to ensure that the modified gene remains within the same translational reading frame as Nr-CAM, uninterrupted by translational stop signals, in the gene region where the desired Nr-CAM activity is encoded.

Additionally, the Nr-CAM-encoding nucleic acid sequence can be mutated in vitro or in vivo, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy preexisting ones, to facilitate further in vitro modification. Any technique for mutagenesis known in the art can be used, including but not limited to, chemical mutagenesis, in vitro site-directed mutagenesis (Hutchinson, C., et al., 1978, J. Biol. Chem 253:6551), use of TAB® linkers (Pharmacia), etc.

Manipulations of the Nr-CAM sequence may also be made at the protein level. Included within the scope of the invention are Nr-CAM protein fragments or other derivatives or analogs which are differentially modified during or after translation, e.g., by glycosylation, acetylation,

20 phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH₄; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

In addition, analogs and derivatives of Nr-CAM can be chemically synthesized. For example, a peptide corresponding to a portion of a Nr-CAM protein which comprises the desired domain, or which mediates the desired activity in vitro, can be synthesized by use of a peptide

synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the Nr-CAM sequence. Nonclassical amino acids include but are not limited to the Ds isomers of the common amino acids, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, γ-Abu, ε-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, tbutylglycine, t-butylalanine, phenylglycine, 10 cyclohexylalanine, β-alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, $C\alpha$ -methyl amino acids, No-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

15 In a specific embodiment, the Nr-CAM derivative is a chimeric, or fusion, protein comprising a Nr-CAM protein or fragment thereof (preferably consisting of at least a domain or motif of the Nr-CAM protein, or at least 10 amino acids of the Nr-CAM protein) joined at its amino- or carboxy-terminus 20 via a peptide bond to an amino acid sequence of a different protein. In one embodiment, such a chimeric protein is produced by recombinant expression of a nucleic acid encoding the protein (comprising a Nr-CAM-coding sequence joined inframe to a coding sequence for a different protein). chimeric product can be made by ligating the appropriate 25 nucleic acid sequences encoding the desired amino acid sequences to each other by methods known in the art, in the proper coding frame, and expressing the chimeric product by methods commonly known in the art. Alternatively, such a chimeric product may be made by protein synthetic techniques, 30 e.g., by use of a peptide synthesizer. Chimeric genes

o e.g., by use of a peptide synthesizer. Chimeric genes comprising portions of Nr-CAM fused to any heterologous protein-encoding sequences may be constructed. A specific

embodiment relates to a chimeric protein comprising a fragment of Nr-CAM of at least six amino acids.

In another specific embodiment, the Nr-CAM derivative is a molecule comprising a region of homology with 5 a Nr-CAM protein. By way of example, in various embodiments, a first protein region can be considered "homologous" to a second protein region when the amino acid sequence of the first region is at least 30%, 40%, 50%, 60%, 70%, 75%, 80%, 90%, or 95% identical, when compared to any sequence in the second region of an equal number of amino acids as the number contained in the first region or when compared to an aligned sequence of the second region that has been aligned by a computer homology program known in the art. For example, a molecule can comprise one or more regions homologous to a Nr-CAM domain or a portion thereof.

Other specific embodiments of derivatives and analogs are described in the subsections below and examples sections infra.

5.3. ASSAYS OF Nr-CAM PROTEINS, DERIVATIVES AND ANALOGS

20

The functional activity of Nr-CAM proteins, derivatives and analogs can be assayed by various methods.

For example, in one embodiment, where one is assaying for the ability to bind or compete with wild-type Nr-CAM for binding to anti-Nr-CAM antibody, various immunoassays known in the art can be used, including but not limited to competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g.,

gel agglutination assays, hemagglutination assays),
complement fixation assays, immunofluorescence assays,
protein A assays, and immunoelectrophoresis assays, etc. In
one embodiment, antibody binding is detected by detecting a
label on the primary antibody. In another embodiment, the
primary antibody is detected by detecting binding of a
secondary antibody or reagent to the primary antibody. In a
further embodiment, the secondary antibody is labelled. Many
means are known in the art for detecting binding in an
immunoassay and are within the scope of the present
invention.

In another embodiment, where a Nr-CAM-binding protein is identified, the binding can be assayed, e.g., by means well-known in the art. In another embodiment, physiological correlates of Nr-CAM binding to its substrates (signal transduction) can be assayed.

In addition, assays that can be used to detect or measure the ability to inhibit, or alternatively promote, cell proliferation are described in Section 5.4.

Other methods will be known to the skilled artisan and are within the scope of the invention.

5.4. DIAGNOSIS AND SCREENING

Nr-CAM proteins, analogs, derivatives, and subsequences thereof, Nr-CAM nucleic acids (and sequences complementary thereto), anti-Nr-CAM antibodies, have uses in diagnostics. Such molecules can be used in assays, such as immunoassays, to detect, prognose, diagnose, or monitor various conditions, diseases, and disorders affecting Nr-CAM expression, or monitor the treatment thereof. In particular, such an immunoassay is carried out by a method comprising contacting a sample derived from a patient with an anti-Nr-CAM antibody under conditions such that immunospecific binding can occur, and detecting or measuring the amount of

any immunospecific binding by the antibody. In a specific aspect, such binding of antibody, in tissue sections, can be used to detect aberrant Nr-CAM localization or aberrant (e.g., high, low or absent) levels of Nr-CAM. In a specific 5 embodiment, antibody to Nr-CAM can be used to assay in a patient tissue or serum sample for the presence of Nr-CAM where an aberrant level of Nr-CAM is an indication of a diseased condition. By "aberrant levels," is meant increased or decreased levels relative to that present, or a standard level representing that present, in an analogous sample from a portion of the body or from a subject not having the disorder. In a specific embodiment, antibody to Nr-CAM can be used to assay and screen tissues or bodily fluids including but not limited to spinal fluid and brain tissue for elevated levels of Nr-CAM expression indicative of a 15 tumor.

The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few.

Nr-CAM genes and related nucleic acid sequences and subsequences, including complementary sequences, can also be used in hybridization assays. Nr-CAM nucleic acid sequences, or subsequences thereof comprising about at least 8 nucleotides, can be used as hybridization probes.

Hybridization assays can be used to detect, prognose, diagnose, or monitor conditions, disorders, or disease states associated with aberrant changes in Nr-CAM expression and/or activity as described supra. In particular, such a

hybridization assay is carried out by a method comprising contacting a sample containing nucleic acid with a nucleic acid probe capable of hybridizing to Nr-CAM DNA or RNA, under conditions such that hybridization can occur, and detecting or measuring any resulting hybridization.

In specific embodiments, diseases and disorders involving overproliferation of cells can be diagnosed, or their suspected presence can be screened for, or a predisposition to develop such disorders can be detected, by detecting increased levels of Nr-CAM protein, Nr-CAM RNA, or 10 Nr-CAM functional activity or by detecting mutations in Nr-CAM RNA, DNA or protein (e.q., translocations in Nr-CAM nucleic acids, truncations in the Nr-CAM gene or protein, changes in nucleotide or amino acid sequence relative to wild-type Nr-CAM) that cause increased expression or activity 15 of Nr-CAM. Such diseases and disorders include but are not limited to those tumors or tissue types mentioned in Section 6 in which Nr-CAM is overexpressed. By way of example, levels of Nr-CAM protein can be detected by immunoassay, levels of Nr-CAM RNA can be detected by hybridization assays 20 (e.g., Northern blots, dot blots), translocations and point mutations in Nr-CAM nucleic acids can be detected by Southern blotting, RFLP analysis, PCR using primers that preferably generate a fragment spanning at least most of the Nr-CAM gene, sequencing of the Nr-CAM genomic DNA or cDNA obtained from the patient, etc. 25

In a preferred embodiment, levels of Nr-CAM mRNA or protein in a patient sample are detected or measured, in which increased levels indicate that the subject has, or has a predisposition to developing, a malignancy or hyperproliferative disorder; in which the increased levels are relative to the levels present in an analogous sample from a portion of the body or from a subject not having the

malignancy or hyperproliferative disorder, as the case may be.

In another specific embodiment, diseases and disorders involving a deficiency in cell proliferation or in s which cell proliferation is desirable for treatment, are diagnosed, or their suspected presence can be screened for, or a predisposition to develop such disorders can be detected, by detecting decreased levels of Nr-CAM protein, Nr-CAM RNA, or Nr-CAM functional activity, or by detecting mutations in Nr-CAM RNA, DNA or protein (e.g., translocations in Nr-CAM nucleic acids, truncations in the gene or protein, changes in nucleotide or amino acid sequence relative to wild-type Nr-CAM) that cause decreased expression or activity of Nr-CAM. Such diseases and disorders include but are not limited to those tumors and tissue types mentioned in Section 15 6 and its subsections in which Nr-CAM is overexpressed. way of example, levels of Nr-CAM protein, levels of Nr-CAM RNA, Nr-CAM binding activity, and the presence of translocations or point mutations can be determined as described above.

In a specific embodiment, levels of Nr-CAM mRNA or protein in a patient sample are detected or measured, in which decreased levels indicate that the subject has, or has a predisposition to developing, a malignancy or hyperproliferative disorder; in which the decreased levels are relative to the levels present in an analogous sample from a portion of the body or from a subject not having the malignancy or hyperproliferative disorder, as the case may be.

Kits for diagnostic use are also provided, that comprise, in one or more containers, an anti-Nr-CAM antibody, and, optionally, a labeled binding partner to the antibody. Alternatively, the anti-Nr-CAM antibody can be labeled (with a detectable marker, e.g., a chemiluminescent, enzymatic,

fluorescent, or radioactive moiety). A kit is also provided that comprises, in one or more containers, a nucleic acid probe capable of hybridizing to Nr-CAM RNA. In a specific embodiment, a kit can comprise in one or more containers a pair of primers (e.g., each in the size range of 6-30 nucleotides) that are capable of priming amplification [e.g., by polymerase chain reaction (see e.g., Innis et al., 1990, PCR Protocols, Academic Press, Inc., San Diego, CA), ligase chain reaction (see EP 320,308) use of Qβ replicase, cyclic probe reaction, or other methods known in the art] under appropriate reaction conditions of at least a portion of a Nr-CAM nucleic acid. A kit can optionally further comprise, in a container, a predetermined amount of a purified Nr-CAM protein or nucleic acid, e.g., for use as a standard or control.

15

5.5. THERAPEUTIC USES

The invention provides for treatment, inhibition or prevention of various diseases and disorders by administration of a therapeutic compound (termed herein 20 "Therapeutic"). Such "Therapeutics" include but are not limited to: Nr-CAM proteins and analogs and derivatives (including fragments) thereof (e.g., as described hereinabove); antibodies thereto (as described hereinabove); nucleic acids encoding the Nr-CAM proteins, analogs, or derivatives (e.g., as described hereinabove); Nr-CAM antisense nucleic acids, and Nr-CAM agonists and antagonists. Disorders involving tumorigenesis or cell overproliferation are treated or prevented by administration of a Therapeutic that antagonizes Nr-CAM function. Disorders in which cell proliferation is deficient or is desired are treated or prevented by administration of a Therapeutic that promotes

Nr-CAM function. See details in the subsections below.

Generally, it is preferred to administer a product of a species origin or species reactivity (in the case of antibodies) that is the same as that of the recipient. Thus, in a preferred embodiment, a human Nr-CAM protein, derivative, or analog, or nucleic acid, or an antibody to a human Nr-CAM protein, is therapeutically or prophylactically administered to a human patient.

Additional descriptions and sources of Therapeutics that can be used according to the invention are found in Sections 5.1 through 5.7 herein.

5.5.1. TREATMENT, INHIBITION AND PREVENTION OF DISORDERS INVOLVING OVERPROLIFERATION OF CELLS

Diseases and disorders involving cell

overproliferation are treated, inhibited or prevented by
administration of a Therapeutic that antagonizes (i.e.,
inhibits) Nr-CAM function. Examples of such a Therapeutic
include but are not limited to Nr-CAM antibodies, Nr-CAM
antisense nucleic acids, derivatives, or analogs that are
functionally active, particularly that are active in
inhibiting cell proliferation (e.g., as demonstrated in in
vitro assays or in animal models or in Drosophila). Other
Therapeutics that can be used, e.g., Nr-CAM antagonists, can
be identified using in vitro assays or animal models,
examples of which are described infra.

In specific embodiments, Therapeutics that inhibit Nr-CAM function are administered therapeutically (including prophylactically): (1) in diseases or disorders involving an increased (relative to normal or desired) level of Nr-CAM protein or function, for example, in patients where Nr-CAM protein is overexpressed, genetically defective, or biologically hyperactive; or (2) in diseases or disorders wherein in vitro (or in vivo) assays (see infra) indicate the

utility of Nr-CAM antagonist administration. The increased level in Nr-CAM protein or function can be readily detected, e.g., by obtaining a patient tissue sample (e.g., from biopsy tissue) and assaying it in vitro for RNA or protein levels, structure and/or activity of the expressed Nr-CAM RNA or protein. Many methods standard in the art can be thus employed, including but not limited to immunoassays to detect and/or visualize Nr-CAM protein (e.g., Western blot, immunoprecipitation followed by sodium dodecyl sulfate polyacrylamide gel electrophoresis, immunocytochemistry, etc.) and/or hybridization assays to detect Nr-CAM expression by detecting and/or visualizing Nr-CAM mRNA (e.g., Northern assays, dot blots, in situ hybridization, etc.), etc.

Diseases and disorders involving cell overproliferation that can be treated, inhibited or prevented include but are not limited to malignancies, premalignant conditions (e.g., hyperplasia, metaplasia, dysplasia), benign tumors, hyperproliferative disorders, benign dysproliferative disorders, etc. Examples of these are detailed below.

5.5.1.1. MALIGNANCIES

Malignancies and related disorders that can be treated or prevented by administration of a Therapeutic that inhibits Nr-CAM function include but are not limited to those listed in Table 1 (for a review of such disorders, see

Fishman et al., 1985, Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia).

30

20

TABLE 1 MALIGNANCIES AND RELATED DISORDERS

Leukemia acute leukemia 5 acute lymphocytic leukemia acute lymphoblastic leukemia acute myelocytic leukemia myeloblastic myelogenous promyelocytic myelomonocytic 10 monocytic erythroleukemia chronic leukemia chronic myelocytic (granulocytic) leukemia chronic myelogenous leukemia chronic lymphocytic leukemia Polycythemia vera Lymphoma Hodgkin's disease 15 non-Hodgkin's disease Multiple myeloma Waldenström's macroglobulinemia Heavy chain disease Solid tumors sarcomas and carcinomas adenocarcinoma 20 fibrosarcoma myxosarcoma liposarcoma chondrosarcoma osteogenic sarcoma chordoma angiosarcoma endotheliosarcoma lymphangiosarcoma 25 lymphangioendotheliosarcoma synovioma mesothelioma Ewing's tumor leiomyosarcoma rhabdomyosarcoma colon carcinoma colorectal adenocarcinoma 30 colon tumor metastatic to brain lung carcinoma pancreatic cancer breast cancer

ovarian cancer prostate cancer squamous cell carcinoma basal cell carcinoma adenocarcinoma sweat gland carcinoma sebaceous gland carcinoma 5 papillary carcinoma papillary adenocarcinomas cystadenocarcinoma medullary carcinoma bronchogenic carcinoma renal cell carcinoma hepatoma bile duct carcinoma 10 choriocarcinoma seminoma embryonal carcinoma Wilms' tumor cervical cancer uterine cancer testicular tumor lung carcinoma 15 small cell lung carcinoma bladder carcinoma epithelial carcinoma glioblastoma glioma astrocytoma medulloblastoma craniopharyngioma 20 ependymoma pinealoma hemangioblastoma acoustic neuroma oligodendroglioma meningioma melanoma neuroblastoma 25 retinoblastoma

In specific embodiments, malignancy or dysproliferative changes (such as metaplasias and dysplasias), or hyperproliferative disorders, are treated, inhibited or prevented in the brain. In other specific

embodiments, carcinoma, melanoma, or leukemia is treated, inhibited or prevented.

5.5.1.2. PREMALIGNANT CONDITIONS

- Nr-CAM activity can also be administered to treat, inhibit premalignant conditions and to inhibit or prevent progression to a neoplastic or malignant state, including but not limited to those disorders listed in Table 1. Such prophylactic or therapeutic use is indicated in conditions known or suspected of preceding progression to neoplasia or cancer, in particular, where non-neoplastic cell growth consisting of hyperplasia, metaplasia, or most particularly, dysplasia has occurred (for review of such abnormal growth conditions, see Robbins and Angell, 1976, Basic Pathology, 2d Ed., W.B.
- 15 Saunders Co., Philadelphia, pp. 68-79.) Hyperplasia is a form of controlled cell proliferation involving an increase in cell number in a tissue or organ, without significant alteration in structure or function. As but one example, endometrial hyperplasia often precedes endometrial cancer.
- 20 Metaplasia is a form of controlled cell growth in which one type of adult or fully differentiated cell substitutes for another type of adult cell. Metaplasia can occur in epithelial or connective tissue cells. Atypical metaplasia involves a somewhat disorderly metaplastic epithelium.
- Dysplasia is frequently a forerunner of cancer, and is found mainly in the epithelia; it is the most disorderly form of non-neoplastic cell growth, involving a loss in individual cell uniformity and in the architectural orientation of cells. Dysplastic cells often have abnormally large, deeply stained nuclei, and exhibit pleomorphism. Dysplasia
- 30 characteristically occurs where there exists chronic irritation or inflammation, and is often found in the cervix, respiratory passages, oral cavity, and gall bladder.

Alternatively or in addition to the presence of abnormal cell growth characterized as hyperplasia, metaplasia, or dysplasia, the presence of one or more characteristics of a transformed phenotype, or of a malignant s phenotype, displayed in vivo or displayed in vitro by a cell sample from a patient, can indicate the desirability of prophylactic/therapeutic administration of a Therapeutic that inhibits Nr-CAM function. As mentioned supra, such characteristics of a transformed phenotype include morphology changes, looser substratum attachment, loss of contact inhibition, loss of anchorage dependence, protease release, increased sugar transport, decreased serum requirement, expression of fetal antigens, disappearance of the 250,000 dalton cell surface protein, etc. (see also id., at pp. 84-90 for characteristics associated with a transformed or 15 malignant phenotype).

In a specific embodiment, leukoplakia, a benignappearing hyperplastic or dysplastic lesion of the epithelium, or Bowen's disease, a carcinoma in situ, are preneoplastic lesions indicative of the desirability of 20 prophylactic intervention.

In another embodiment, fibrocystic disease (cystic hyperplasia, mammary dysplasia, particularly adenosis (benign epithelial hyperplasia)) is indicative of the desirability of prophylactic intervention.

or more of the following predisposing factors for malignancy is treated by administration of an effective amount of a Therapeutic: a chromosomal translocation associated with a malignancy (e.g., the Philadelphia chromosome for chronic myelogenous leukemia, t(14;18) for follicular lymphoma, etc.), familial polyposis or Gardner's syndrome (possible forerunners of colon cancer), benign monoclonal gammopathy (a possible forerunner of multiple myeloma), and a first degree

kinship with persons having a cancer or precancerous disease showing a Mendelian (genetic) inheritance pattern (e.g., familial polyposis of the colon, Gardner's syndrome, hereditary exostosis, polyendocrine adenomatosis, medullary 5 thyroid carcinoma with amyloid production and pheochromocytoma, Peutz-Jeghers syndrome, neurofibromatosis of Von Recklinghausen, retinoblastoma, carotid body tumor, cutaneous melanocarcinoma, intraocular melanocarcinoma, xeroderma pigmentosum, ataxia telangiectasia, Chediak-Higashi syndrome, albinism, Fanconi's aplastic anemia, and Bloom's syndrome; see Robbins and Angell, 1976, Basic Pathology, 2d Ed., W.B. Saunders Co., Philadelphia, pp. 112-113) etc.) In another specific embodiment, a Therapeutic of the invention is administered to a human patient to prevent progression to brain, breast, colon, prostate, lung, or skin. 15 In other specific embodiments, carcinoma, melanoma, or leukemia is treated or prevented.

5.5.1.3. GENE THERAPY

In a specific embodiment, anti-sense nucleic acids

20 complementary to a sequence encoding a Nr-CAM protein or
functional derivative thereof, are administered to inhibit
Nr-CAM function, by way of gene therapy. Gene therapy refers
to therapy performed by the administration of a nucleic acid
to a subject. In this embodiment of the invention, the
antisense nucleic acid mediates a therapeutic effect by
inhibiting Nr-CAM transcription and translation.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

For general reviews of the methods of gene therapy, see Goldspiel et al., 1993, Clinical Pharmacy 12:488-505; Wu and Wu, 1991, Biotherapy 3:87-95; Tolstoshev, 1993, Ann. Rev. Pharmacol. Toxicol. 32:573-596; Mulligan, 1993, Science

260:926-932; and Morgan and Anderson, 1993, Ann. Rev.
Biochem. 62:191-217; May, 1993, TIBTECH 11(5):155-215).
Methods commonly known in the art of recombinant DNA
technology which can be used are described in Ausubel et al.,
(eds.), 1993, Current Protocols in Molecular Biology, John
Wiley & Sons, NY; and Kriegler, 1990, Gene Transfer and
Expression, A Laboratory Manual, Stockton Press, NY.

In one embodiment, the Therapeutic comprises an Nr-CAM sense or antisense nucleic acid that is part of an expression vector that expresses a Nr-CAM protein or fragment or chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the Nr-CAM coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, a nucleic acid molecule is used in which the Nr-CAM coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the Nr-CAM nucleic acid (Koller and Smithies, 1989, Proc. Natl. Acad.

Sci. USA 86:8932-8935; Zijlstra et al., 1989, Nature 342:435-

Delivery of the nucleic acid into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid in vitro, then transplanted into the patient.

These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

438).

In a specific embodiment, the nucleic acid is directly administered in vivo, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing it as part of an appropriate nucleic acid expression vector

and administering it so that it becomes intracellular, e.g., by infection using a defective or attenuated retroviral or other viral vector (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle 5 bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see e.g., Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, a nucleic acidligand complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the 15 nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180 dated April 16, 1992 (Wu et al.); WO 92/22635 dated December 23, 1992 20 (Wilson et al.); WO92/20316 dated November 26, 1992 (Findeis et al.); WO93/14188 dated July 22, 1993 (Clarke et al.), WO 93/20221 dated October 14, 1993 (Young)). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by 25 homologous recombination (Koller and Smithies, 1989, Proc. Natl. Acad. Sci. USA 86:8932-8935; Zijlstra et al., 1989, Nature 342:435-438).

In a specific embodiment, a viral vector that contains the Nr-CAM nucleic acid is used. For example, a retroviral vector can be used (see Miller et al., 1993, Meth. Enzymol. 217:581-599; Kondo, et al., 1998, Cancer Res, 68:962-967; Boviatsis, et al., 1994, Human Gene Therapy, 5:183-191. These retroviral vectors have been modified to

delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The Nr-CAM nucleic acid to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., 1994, Biotherapy 6:291-302, which describes the use of a retroviral vector to deliver the

mdr1 gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., 1994, J. Clin. Invest. 93:644-651; Kiem et al., 1994, Blood 83:1467-1473; Salmons and Gunzberg, 1993, Human Gene Therapy 4:129-141; and Grossman and Wilson, 1993, Curr. Opin. in Genetics and Devel. 3:110-114.

Adenoviruses are other viral vectors that can be 15 used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirusbased delivery systems are liver, the central nervous system, 20 endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, 1993, Current Opinion in Genetics and Development 3:499-503 present a review of adenovirus-based qene therapy. Bout et al., 1994, Human Gene Therapy 5:3-10demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., 1991, Science 252:431-434; Rosenfeld et al., 1992, Cell 68:143-155; and Mastrangeli et al., 1993, J. Clin. Invest. 91:225-234.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., 1993, Proc. Soc. Exp. Biol. Med. 204:289-300.

30

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid 15 sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see e.g., Loeffler and Behr, 1993, Meth. Enzymol. 217:599-618; Cohen et al., 1993, Meth. 20 Enzymol. 217:618-644; Cline, 1985, Pharmac. Ther. 29:69-92) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. In a preferred embodiment, epithelial cells are injected, e.g., subcutaneously. In another embodiment, recombinant skin cells may be applied as a skin graft onto the patient. Recombinant blood cells (e.g., hematopoietic stem or

progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils,

eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a preferred embodiment, the cell used for gene 15 therapy is autologous to the patient.

In an embodiment in which recombinant cells are used in gene therapy, a Nr-CAM nucleic acid is introduced into the cells such that it is expressible by the cells or their progeny, and the recombinant cells are then

- administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention. Such stem cells include but are not limited to hematopoietic stem cells (HSC), stem cells of epithelial tissues such as the skin and the lining of the gut, embryonic heart muscle cells, liver stem cells (PCT Publication WO 94/08598, dated April 28, 1994), and
 - neural stem cells (Stemple and Anderson, 1992, Cell 71:973-985).

30

Epithelial stem cells (ESCs) or keratinocytes can be obtained from tissues such as the skin and the lining of the gut by known procedures (Rheinwald, 1980, Meth. Cell Bio.

21A:229). In stratified epithelial tissue such as the skin, renewal occurs by mitosis of stem cells within the germinal layer, the layer closest to the basal lamina. Stem cells within the lining of the gut provide for a rapid renewal rate of this tissue. ESCs or keratinocytes obtained from the skin or lining of the gut of a patient or donor can be grown in tissue culture (Rheinwald, 1980, Meth. Cell Bio. 21A:229; Pittelkow and Scott, 1986, Mayo Clinic Proc. 61:771). If the ESCs are provided by a donor, a method for suppression of host versus graft reactivity (e.g., irradiation, drug or antibody administration to promote moderate immunosuppression) can also be used.

With respect to hematopoietic stem cells (HSC), any technique which provides for the isolation, propagation, and maintenance in vitro of HSC can be used in this embodiment of 15 the invention. Techniques by which this may be accomplished include (a) the isolation and establishment of HSC cultures from bone marrow cells isolated from the future host, or a donor, or (b) the use of previously established long-term HSC cultures, which may be allogeneic or xenogeneic. Non-20 autologous HSC are used preferably in conjunction with a method of suppressing transplantation immune reactions of the future host/patient. In a particular embodiment of the present invention, human bone marrow cells can be obtained from the posterior iliac crest by needle aspiration (see, e.g., Kodo et al., 1984, J. Clin. Invest. 73:1377-1384). In a preferred embodiment of the present invention, the HSCs can be made highly enriched or in substantially pure form. enrichment can be accomplished before, during, or after longterm culturing, and can be done by any techniques known in the art. Long-term cultures of bone marrow cells can be

30 established and maintained by using, for example, modified Dexter cell culture techniques (Dexter et al., 1977, J. Cell

Physiol. <u>91</u>:335) or Witlock-Witte culture techniques (Witlock and Witte, 1982, Proc. Natl. Acad. Sci. USA <u>79</u>:3608-3612).

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

Additional methods that can be adapted for use to deliver a nucleic acid encoding a Nr-CAM protein or functional derivative thereof are described in Section 5.5.2.2.2., infra.

5.5.2. TREATMENT, INHIBITION AND PREVENTION OF HYPERPROLIFERATIVE AND DYSPROLIFERATIVE DISORDERS

15

Diseases and disorders involving an increase in cell proliferation (growth) or in which cell proliferation is otherwise undesirable, are treated, inhibited or prevented by administration of a Therapeutic that antagonizes (inhibits) 20 Nr-CAM function. Therapeutics that can be used include but are not limited to anti-Nr-CAM antibodies (and fragments and derivatives thereof containing the binding region thereof), Nr-CAM antisense nucleic acids, and Nr-CAM nucleic acids that are dysfunctional (e.g., due to a heterologous (non-Nr-CAM 25 sequence) insertion within the Nr-CAM coding sequence) that are used to "knockout" endogenous Nr-CAM function by homologous recombination (see, e.g., Capecchi, 1989, Science 244:1288-1292). In a specific embodiment of the invention, a nucleic acid containing a portion of a Nr-CAM gene in which Nr-CAM sequences flank (are both 5' and 3' to) a different 30 gene sequence, is used, as a Nr-CAM antagonist, to promote Nr-CAM inactivation by homologous recombination (see also Koller and Smithies, 1989, Proc. Natl. Acad. Sci. USA

86:8932-8935; Zijlstra et al., 1989, Nature 342:435-438).
Other Therapeutics that inhibit Nr-CAM function can be
identified by use of known convenient in vitro assays, e.g.,
based on their ability to inhibit binding of Nr-CAM to
sanother protein or inhibit any known Nr-CAM function, as
preferably assayed in vitro or in cell culture, although
genetic assays in Drosophila or another species may also be
employed. Preferably, suitable in vitro or in vivo assays,
are utilized to determine the effect of a specific
Therapeutic and whether its administration is indicated for
treatment of the affected tissue.

In specific embodiments, Therapeutics that inhibit Nr-CAM function are administered therapeutically (including prophylactically): (1) in diseases or disorders involving an increased (relative to normal or desired) level of Nr-CAM 15 protein or function, for example, in patients where Nr-CAM protein is overactive or overexpressed; or (2) in diseases or disorders wherein in vitro (or in vivo) assays (see infra) indicate the utility of Nr-CAM antagonist administration. The increased levels in Nr-CAM protein or function can be 20 readily detected, e.g., by quantifying protein and/or RNA, by obtaining a patient tissue sample (e.g., from biopsy tissue) and assaying it in vitro for RNA or protein levels, structure and/or activity of the expressed Nr-CAM RNA or protein. methods standard in the art can be thus employed, including 25 but not limited to immunoassays to detect and/or visualize Nr-CAM protein (e.g., Western blot, immunoprecipitation followed by sodium dodecyl sulfate polyacrylamide gel electrophoresis, immunocytochemistry, etc.) and/or hybridization assays to detect Nr-CAM expression by detecting and/or visualizing respectively Nr-CAM mRNA (e.g., Northern assays, dot blots, in situ hybridization, etc.), etc.

In other embodiments, chemical mutagenesis, or homologous recombination with an insertionally inactivated

Nr-CAM gene (see Capecchi, 1989, Science 244:1288-1292 and Section 5.14 infra) can be carried out to reduce or destroy endogenous Nr-CAM function, in order to decrease cell proliferation. Suitable methods, modes of administration and compositions, that can be used to inhibit Nr-CAM function are described in Sections 5.8.2 through 5.8.2.1.2, above.

In an embodiment of the invention, a Therapeutic that inhibits Nr-CAM activity is used to treat, inhibit or prevent hyperproliferative or benign dysproliferative disorders. Specific embodiments are directed to treatment, inhibition or prevention of cirrhosis of the liver (a condition in which scarring has overtaken normal liver regeneration processes), treatment or inhibition of keloid (hypertrophic scar) formation (disfiguring of the skin in which the scarring process interferes with normal renewal), psoriasis (a common skin condition characterized by excessive proliferation of the skin and delay in proper cell fate determination), benign tumors, fibrocystic conditions, and tissue hypertrophy (e.g., prostatic hyperplasia).

20 5.5.2.1. ANTISENSE REGULATION OF Nr-CAM EXPRESSION

In a specific embodiment, Nr-CAM function is inhibited by use of Nr-CAM antisense nucleic acids. The present invention provides the therapeutic or prophylactic use of nucleic acids of at least six nucleotides that are antisense to a gene or cDNA encoding Nr-CAM or a portion thereof. A Nr-CAM "antisense" nucleic acid as used herein refers to a nucleic acid capable of hybridizing to a portion of a Nr-CAM RNA (preferably mRNA) by virtue of some sequence complementarity. The antisense nucleic acid may be complementary to a coding and/or noncoding region of a Nr-CAM mRNA. Such antisense nucleic acids have utility as

Therapeutics that inhibits Nr-CAM function, and can be used

in the treatment or prevention of disorders as described supra in Section 5.5.2 and its subsections.

The antisense nucleic acids of the invention can be oligonucleotides that are double-stranded or single-stranded, 5 RNA or DNA or a modification or derivative thereof, which can be directly administered to a cell, or which can be produced intracellularly by transcription of exogenous, introduced sequences.

In a specific embodiment, the Nr-CAM antisense nucleic acids provided by the present invention can be used to inhibit or prevent tumors or other forms of aberrant cell proliferation.

The invention further provides pharmaceutical compositions comprising an effective amount of the Nr-CAM antisense nucleic acids of the invention in a

15 pharmaceutically acceptable carrier, as described infra.

In another embodiment, the invention is directed to methods for inhibiting the expression of a Nr-CAM nucleic acid sequence in a prokaryotic or eukaryotic cell comprising providing the cell with an effective amount of a composition comprising an Nr-CAM antisense nucleic acid of the invention.

Nr-CAM antisense nucleic acids and their uses are described in detail below.

5.5.2.1.1. Nr-CAM ANTISENSE NUCLEIC ACIDS

The Nr-CAM antisense nucleic acids are of at least six nucleotides and are preferably oligonucleotides (ranging from 6 to about 50 oligonucleotides). In specific aspects, the oligonucleotide is at least 10 nucleotides, at least 15 nucleotides, at least 100 nucleotides, or at least 200 nucleotides. The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. In a specific embodiment, the antisense nucleic acids of the invention are

double-stranded RNA (see, Fire et al., 1998, Nature 391:806-811). The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO 88/09810, published December 15, 1988) or blood-brain barrier (see, e.g., PCT Publication No. WO 89/10134, published April 25, 1988), hybridization-triggered cleavage agents (see, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents (see, e.g., Zon, 1988, Pharm. Res. 5:539-549).

In a preferred aspect of the invention, a Nr-CAM antisense oligonucleotide is provided, preferably of single-stranded DNA. The oligonucleotide may be modified at any position on its structure with substituents generally known in the art.

The Nr-CAM antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxylmethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil,

queosine, 2-thiocytosine, 5-methyl-2-thiouracil,

2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

In another embodiment, the oligonucleotide comprises at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothicate, a phosphorodithicate, a phosphoramidate, a phosphoramidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641).

The oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

oligonucleotides of the invention may be
synthesized by standard methods known in the art, e.g. by use
of an automated DNA synthesizer (such as are commercially
available from Biosearch, Applied Biosystems, etc.). As
examples, phosphorothicate oligonucleotides may be
synthesized by the method of Stein et al. (1988, Nucl. Acids
Res. 16:3209), methylphosphonate oligonucleotides can be
prepared by use of controlled pore glass polymer supports
(Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:74487451), etc.

In a specific embodiment, the Nr-CAM antisense oligonucleotide comprises catalytic RNA, or a ribozyme (see, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al., 1990, Science 247:1222-1225).

5 In another embodiment, the oligonucleotide is a 2'-0-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analog (Inoue et al.,

In an alternative embodiment, the Nr-CAM antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector can be introduced in vivo such that it is taken up by a cell, within which cell the vector or a portion thereof is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the Nr-CAM antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and

20 or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the Nr-CAM antisense RNA can be by any promoter known in the art to act in mammalian, preferably human, cells. Such promoters can be inducible or constitutive.

Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, Nature 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, Cell 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445), the regulatory

sequences of the metallothionein gene (Brinster et al., 1982,

Nature 296:39-42), etc.

1987, FEBS Lett. 215:327-330).

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a Nr-CAM nucleic acid or Nr-CAM gene, preferably a human Nr-CAM gene. However, absolute s complementarily, although preferred, is not required. sequence "complementary to at least a portion of an RNA," as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded Nr-CAM antisense nucleic acids, a single strand of the duplex DNA 10 may thus be tested, or triplex formation may be assayed. ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with a Nr-CAM RNA it may contain and still 15 form a stable duplex (or triplex, as the case may be). skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

20 5.5.2.1.2. THERAPEUTIC USE OF Nr-CAM ANTISENSE NUCLEIC ACIDS

The Nr-CAM antisense nucleic acids can be used to treat, inhibit (or prevent) disorders of a cell type that expresses, or preferably overexpresses, Nr-CAM. In a specific embodiment, such a disorder is a hyperproliferative disorder, e.g. tumorigenesis. In a preferred embodiment, a single-stranded DNA antisense Nr-CAM oligonucleotide is used.

Cell types which express or overexpress Nr-CAM RNA can be identified by various methods known in the art. Such methods include but are not limited to hybridization with a 30 Nr-CAM-specific nucleic acid (e.g. by Northern hybridization, dot blot hybridization, in situ hybridization), observing the ability of RNA from the cell type to be translated in vitro

into Nr-CAM, immunoassay, etc. In a preferred aspect, primary tissue from a patient can be assayed for Nr-CAM expression prior to treatment, e.g., by immunocytochemistry or in situ hybridization.

5 Pharmaceutical compositions of the invention (see Section 5.10), comprising an effective amount of a Nr-CAM antisense nucleic acid in a pharmaceutically acceptable carrier, can be administered to a patient having a disease or disorder which is of a type that expresses or overexpresses Nr-CAM RNA or protein.

The amount of Nr-CAM antisense nucleic acid which will be effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. Where possible, it is desirable to determine the antisense cytotoxicity of the tumor type to be treated in vitro, and then in useful animal model systems prior to testing and use in humans.

In a specific embodiment, pharmaceutical compositions comprising Nr-CAM antisense nucleic acids are administered via liposomes, microparticles, or microcapsules. In various embodiments of the invention, it may be useful to use such compositions to achieve sustained release of the Nr-CAM antisense nucleic acids. In a specific embodiment, it may be desirable to utilize liposomes targeted via antibodies to specific identifiable tumor antigens (Leonetti et al., 1990, Proc. Natl. Acad. Sci. U.S.A. 87:2448-2451; Renneisen et al., 1990, J. Biol. Chem. 265:16337-16342).

Additional methods that can be adapted for use to deliver a Nr-CAM antisense nucleic acid are described in Section 5.9.1.4.

5.6. DEMONSTRATION OF THERAPEUTIC OR PROPHYLACTIC UTILITY

The Therapeutics of the invention are preferably tested in vitro, and then in vivo for the desired therapeutic or prophylactic activity, prior to use in humans.

For example, in vitro assays which can be used to determine whether administration of a specific Therapeutic is indicated, include in vitro cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a Therapeutic, and the effect of such 10 Therapeutic upon the tissue sample is observed. embodiment, where the patient has a malignancy, a sample of cells from such malignancy is plated out or grown in culture, and the cells are then exposed to a Therapeutic. A Therapeutic which inhibits survival or growth of the 15 malignant cells is selected for therapeutic use in vivo. Many assays standard in the art can be used to assess such survival and/or growth; for example, cell proliferation can be assayed by measuring ³H-thymidine incorporation, by direct cell count, by detecting changes in transcriptional activity 20 of known genes such as proto-oncogenes (e.g., fos, myc) or cell cycle markers; cell viability can be assessed by trypan blue staining, differentiation can be assessed visually based on changes in morphology, etc.

In another embodiment, a Therapeutic is indicated for use which exhibits the desired effect, inhibition or promotion of cell growth, upon a patient cell sample from tissue having or suspected of having a hyper- or hypoproliferative disorder, respectively. Such hyper- or hypoproliferative disorders include but are not limited to those described herein.

In another specific embodiment, a Therapeutic is indicated for use in treating or inhibiting cell injury or a

degenerative disorder which exhibits in vitro promotion of growth/proliferation of cells of the affected patient type.

In various specific embodiments, in vitro assays can be carried out with representative cells of cell types involved in a patient's disorder, to determine if a Therapeutic has a desired effect upon such cell types.

In another embodiment, cells of a patient tissue sample suspected of being pre-neoplastic are similarly plated out or grown in vitro, and exposed to a Therapeutic. Therapeutic which results in a cell phenotype that is more 10 normal (i.e., less representative of a pre-neoplastic state, neoplastic state, malignant state, or transformed phenotype) is selected for therapeutic use. Many assays standard in the art can be used to assess whether a pre-neoplastic state, neoplastic state, or a transformed or malignant phenotype, is 15 present. For example, characteristics associated with a transformed phenotype (a set of in vitro characteristics associated with a tumorigenic ability in vivo) include a more rounded cell morphology, looser substratum attachment, loss of contact inhibition, loss of anchorage dependence, release 20 of proteases such as plasminogen activator, increased sugar transport, decreased serum requirement, expression of fetal antigens, disappearance of the 250,000 dalton surface protein, etc. (see Luria et al., 1978, General Virology, 3d Ed., John Wiley & Sons, New York pp. 436-446).

In other specific embodiments, the in vitro assays described supra can be carried out using a cell line, rather than a cell sample derived from the specific patient to be treated, in which the cell line is derived from or displays characteristic(s) associated with the malignant, neoplastic or pre-neoplastic disorder desired to be treated or prevented, or is derived from the cell type upon which an effect is desired, according to the present invention.

Compounds for use in therapy can be tested in suitable animal model systems prior to testing in humans, including but not limited to rats, mice, chicken, cows, monkeys, rabbits, etc. For in vivo testing, prior to administration to humans, any animal model system known in the art may be used.

5.7. THERAPEUTIC/PROPHYLACTIC ADMINISTRATION AND COMPOSITIONS

The invention provides methods of treatment (and prophylaxis) by administration to a subject of an effective amount of a Therapeutic of the invention. In a preferred aspect, the Therapeutic is isolated, purified or substantially purified. The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human. In a specific embodiment, a non-human mammal is the subject.

Formulations and methods of administration that can be employed when the Therapeutic comprises a nucleic acid are described above; additional appropriate formulations and routes of administration can be selected from among those described hereinbelow.

Various delivery systems are known and can be used to administer a Therapeutic of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the Therapeutic, receptor-mediated endocytosis (see, e.g., Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432), construction of a Therapeutic nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds may be administered by any convenient route, for

example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents.

5 Administration can be systemic or local.

In addition, it may be desirable to introduce a Therapeutic of the invention into the central nervous system by any suitable route, including, but not limited to intraventricular and intrathecal injection. Intraventricular injection may be facilitated by an intraventricular catheter, 10 for example, attached to a reservoir, such as an Ommaya reservoir. Agents which enhance the delivery of chemotherapeutics to brain tumors, such as agonists which activate specific receptors on endothelial cells which regulate permeability, including, e.g., bradykinin agonists 15 (see, e.g., Elliott, et al., 1996, Cancer Research <u>56</u>:3998-4005) tumor angiogenesis factors (Cserr and Knopf, 1992, Immunol Today 12:507-512) etc. can be used in formulations and methods of administration when the Therapeutic is intended for delivery to a tumor of the central nervous 20 system.

In a specific embodiment, injection into spinal fluid, and/or procedures utilizing an Ommaya reservoir, can be used to introduce a therapeutic of the invention such as an anti-Nr-CAM antibody, e.g. a bispecific anti-Nr-CAM antibody, directly into the central nervous system for immunotherapy of a tumor.

25

30

In yet another specific embodiment, an anti-Nr-CAM antibody, e.g. a bispecific anti-Nr-CAM antibody, is employed as a Therapeutic in an immunotherapeutic treatment of a non-brain tumor and is infused into a recipient intravenously.

Immune cells, e.g. dendritic cells or cytotoxic T-cells, can cross the blood-brain barrier and have access to brain tissue, especially in the presence of tumor

angiogenesis factors (Cserr and Knopf, 1992, Immunol. Today, 12:507-512). In a preferred embodiment, activated dendritic cells (HLA-matched to the recipient) (see generally, Tjoa et al., 1996, Prostate 28:65-69) that have been exposed to a Nr-5 CAM protein, analog or derivative thereof are infused into a recipient under conditions that permit their crossing the blood-brain barrier, e.g. in the presence of tumor angiogenesis factors. In another preferred embodiment, activated cytotoxic T-cells (HLA-matched to the recipient) (see generally, Tjoa et al., 1996, Prostate 28:65-69) that have been exposed ex vivo (i.e. in vitro) to a Nr-CAM protein, analog, or derivative thereof are infused into a recipient under conditions that permit their crossing the blood-brain barrier.

In yet another specific embodiment, a Therapeutic of the invention; e.g., activated dendritic cells that have been exposed to a Nr-CAM protein, analog or derivative thereof, or activated cytotoxic T-cells that have been exposed ex vivo dendritic cells that have been exposed to a Nr-CAM protein, analog, or derivative thereof, is administered for the treatment of a non-brain tumor.

Pulmonary administration of a Therapeutic can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to
administer the Therapeutic of the invention locally to the
area in need of treatment; this may be achieved by, for
example, and not by way of limitation, local infusion during
surgery, topical application, e.g., in conjunction with a
wound dressing after surgery, by injection, by means of a
catheter, by means of a suppository, or by means of an
implant, said implant being of a porous, non-porous, or
gelatinous material, including membranes, such as sialastic
membranes, or fibers. In one embodiment, administration can

be by direct injection at the site (or former site) of a malignant tumor or neoplastic or pre-neoplastic tissue.

In another embodiment, the Therapeutic can be delivered in a vesicle, in particular a liposome (see Langer, 1990 Science 249:1527-1533; Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, ibid., pp. 317-327; see generally ibid.)

In yet another embodiment, the Therapeutic can be

delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, supra; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)). In another embodiment, polymeric materials can be used (see Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, J. Macromol. Sci. Rev. Macromol. Chem. 23:61; see also Levy et al., 1985 Science 228:190; During et al.,

20 1989 Ann. Neurol. <u>25</u>:351; Howard et al., 1989 J. Neurosurg. <u>71</u>:105). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, supra, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

In a specific embodiment where the Therapeutic is a nucleic acid encoding a protein Therapeutic, the nucleic acid can be administered in vivo to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector

(see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see e.g., Joliot et al., 1991, Proc. Natl. Acad. Sci. USA 88:1864-1868), etc. Alternatively, a nucleic acid Therapeutic can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

10 The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a Therapeutic, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory 15 agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such 20 pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and 25 glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium 30 chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying

agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the The composition can be formulated as a suppository, 5 with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. 10 Martin. Such compositions will contain a therapeutically effective amount of the Therapeutic, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

15 In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic 20 aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition 30 is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients

may be mixed prior to administration.

The Therapeutics of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the Therapeutic of the invention 10 which will be effective in the treatment of a particular disorder or condition will depend on the nature of the disorder or condition, and can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage 15 ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. However, suitable dosage ranges for 20 intravenous administration are generally about 20-500 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight to 1 mg/kg body weight. Effective doses may be extrapolated from doseresponse curves derived from in vitro or animal model test 25

Suppositories generally contain active ingredient in the range of 0.5% to 10% by weight; oral formulations preferably contain 10% to 95% active ingredient.

systems.

The invention also provides a pharmaceutical pack

or kit comprising one or more containers filled with one or
more of the ingredients of the pharmaceutical compositions of
the invention. Optionally associated with such container(s)

can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

5.7.1. TREATMENT AND PREVENTION OF HYPOPROLIFERATIVE DISORDERS

proliferation or in which cell proliferation is desired for treatment or prevention, and that can be treated or prevented by promoting Nr-CAM function, include but are not limited to degenerative disorders, growth deficiencies, hypoproliferative disorders, physical trauma, lesions, and wounds; for example, to promote wound healing, or to promote regeneration in degenerated, lesioned or injured tissues, etc. In a specific embodiment, nervous system disorders are treated. In another specific embodiment, a disorder that is not of the nervous system is treated.

Lesions which may be treated according to the 20 present invention include but are not limited to the following lesions:

25

30

- (i) traumatic lesions, including lesions caused by physical injury or associated with surgery;
- (ii) ischemic lesions, in which a lack of oxygen results in cell injury or death, e.g., myocardial or cerebral infarction or ischemia, or spinal cord infarction or ischemia;
- (iii) malignant lesions, in which cells are
 destroyed or injured by malignant tissue;
- (iv) infectious lesions, in which tissue is destroyed or injured as a result of infection,

for example, by an abscess or associated with infection by human immunodeficiency virus,

herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis;

- (v) degenerative lesions, in which tissue is destroyed or injured as a result of a degenerative process, including but not limited to nervous system degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis;
- 10 lesions associated with nutritional diseases or disorders, in which tissue is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration;

5

20

25

30

- (vii) lesions associated with systemic diseases including but not limited to diabetes or systemic lupus erythematosus;
- - (ix) demyelinated lesions of the nervous system, in which a portion of the nervous system is destroyed or injured by a demyelinating disease including but not limited to multiple sclerosis, human immunodeficiency virusassociated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients)

according to the invention include but are not limited to the lesions of either the central (including spinal cord, brain) or peripheral nervous systems.

Therapeutics which are useful according to this

5 embodiment of the invention for treatment of a disorder may
be selected by testing for biological activity in promoting
the survival or differentiation of cells (see also Section
5.9). For example, in a specific embodiment relating to
therapy of the nervous system, a Therapeutic which elicits
one of the following effects may be useful according to the
invention:

(i) increased sprouting of neurons in culture or in vivo;

15

- (ii) increased production of a neuron-associated molecule in culture or in vivo, e.g., choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or
- (iii) decreased symptoms of neuron dysfunction in vivo.

Such effects may be measured by any method known in the art.

20 In preferred, non-limiting embodiments, increased sprouting of neurons may be detected by methods set forth in Pestronk et al. (1980, Exp. Neurol. 70:65-82) or Brown et al. (1981, Ann. Rev. Neurosci. 4:17-42); and increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, etc., depending on the molecule to be measured.

5.8. ADDITIONAL USE OF INCREASED Nr-CAM FUNCTION TO PROMOTE INCREASED GROWTH

Promotion of Nr-CAM function (e.g., by

30 administering a compound that promotes Nr-CAM function as

described above), has utility that is not limited to

therapeutic or prophylactic applications. For example, Nr-

CAM function can be promoted in order to increase growth of animals (e.g., cows, horses, pigs, goats, deer, chickens) and plants (particularly edible plants, e.g., tomatoes, melons, lettuce, carrots, potatoes, and other vegetables),

- 5 particularly those that are food or material sources. In an embodiment in which a Nr-CAM nucleic acid is under the control of a tissue-specific promoter, the invention can be used in plants or animals to increase growth where desired (e.g., in the fruit or muscle). For example, a Nr-CAM
- nucleic acid under the control of a temperature-sensitive promoter can be administered to a plant or animal, and the desired portion of the (or the entire) plant or animal can be subjected to heat in order to induce Nr-CAM nucleic acid production, resulting in increased Nr-CAM expression, and resulting cell proliferation. Methods to make plants
- recombinant are commonly known in the art and can be used. Regarding methods of plant transformation (e.g., for transformation with a Nr-CAM antisense nucleic acid), see e.g., Valvekens et al., 1988, Proc. Natl. Acad. Sci. USA 85:5536-5540. Regarding methods of targeted gene
- 20 inactivation in plants (e.g., to inactivate Nr-CAM), see e.g., Miao and Lam, 1995, The Plant J. 7:359-365.

Promotion of Nr-CAM function can also have uses in vitro, e.g., to expand cells in vitro, including but not limited to stem cells, progenitor cells, muscle cells,

25 fibroblasts, liver cells, etc., e.g., to grow cells/tissue in vitro prior to administration to a patient (preferably a patient from which the cells were derived), etc.

5.9. SCREENING FOR Nr-CAM AGONISTS AND ANTAGONISTS

Nr-CAM nucleic acids, proteins, and derivatives also have uses in screening assays to detect molecules that specifically bind to Nr-CAM nucleic acids, proteins, or derivatives and thus have potential use as agonists or

antagonists of Nr-CAM, in particular, molecules that thus affect cell proliferation. In a preferred embodiment, such assays are performed to screen for molecules with potential utility as anti-cancer drugs or lead compounds for drug 5 development. The invention thus provides assays to detect molecules that specifically bind to Nr-CAM nucleic acids, proteins, or derivatives. For example, recombinant cells expressing Nr-CAM nucleic acids can be used to recombinantly produce Nr-CAM proteins in these assays, to screen for molecules that bind to a Nr-CAM protein. Molecules (e.g., 10 putative binding partners of Nr-CAM) are contacted with the Nr-CAM protein (or fragment thereof) under conditions conducive to binding, and then molecules that specifically bind to the Nr-CAM protein are identified. Similar methods can be used to screen for molecules that bind to Nr-CAM 15 derivatives or nucleic acids. Methods that can be used to carry out the foregoing are commonly known in the art.

By way of example, diversity libraries, such as random or combinatorial peptide or nonpeptide libraries can be screened for molecules that specifically bind to Nr-CAM.

20 Many libraries are known in the art that can be used, e.g., chemically synthesized libraries, recombinant (e.g., phage display libraries), and in vitro translation-based libraries.

Examples of chemically synthesized libraries are

described in Fodor et al., 1991, Science 251:767-773;

Houghten et al., 1991, Nature 354:84-86; Lam et al., 1991, Nature 354:82-84; Medynski, 1994, Bio/Technology 12:709-710; Gallop et al., 1994, J. Medicinal Chemistry 37(9):1233-1251; Ohlmeyer et al., 1993, Proc. Natl. Acad. Sci. USA 90:10922-10926; Erb et al., 1994, Proc. Natl. Acad. Sci. USA 91:11422-11426; Houghten et al., 1992, Biotechniques 13:412; Jayawickreme et al., 1994, Proc. Natl. Acad. Sci. USA

91:1614-1618; Salmon et al., 1993, Proc. Natl. Acad. Sci. USA

90:11708-11712; PCT Publication No. WO 93/20242; and Brenner and Lerner, 1992, Proc. Natl. Acad. Sci. USA 89:5381-5383.

Examples of phage display libraries are described in Scott and Smith, 1990, Science 249:386-390; Devlin et al., 1990, Science, 249:404-406; Christian, R.B., et al., 1992, J. Mol. Biol. 227:711-718); Lenstra, 1992, J. Immunol. Meth. 152:149-157; Kay et al., 1993, Gene 128:59-65; and PCT Publication No. WO 94/18318 dated August 18, 1994.

In vitro translation-based libraries include but are not limited to those described in PCT Publication No. WO 91/05058 dated April 18, 1991; and Mattheakis et al., 1994, Proc. Natl. Acad. Sci. USA 91:9022-9026.

By way of examples of nonpeptide libraries, a benzodiazepine library (see e.g., Bunin et al., 1994, Proc. Natl. Acad. Sci. USA 91:4708-4712) can be adapted for use.

Peptoid libraries (Simon et al., 1992, Proc. Natl. Acad. Sci. USA 89:9367-9371) can also be used. Another example of a library that can be used, in which the amide functionalities in peptides have been permethylated to generate a chemically transformed combinatorial library, is described by Ostresh et al. (1994, Proc. Natl. Acad. Sci. USA 91:11138-11142).

Screening the libraries can be accomplished by any of a variety of commonly known methods. See, e.g., the following references, which disclose screening of peptide libraries: Parmley and Smith, 1989, Adv. Exp. Med. Biol.

25 251:215-218; Scott and Smith, 1990, Science 249:386-390; Fowlkes et al., 1992; BioTechniques 13:422-427; Oldenburg et al., 1992, Proc. Natl. Acad. Sci. USA 89:5393-5397; Yu et al., 1994, Cell 76:933-945; Staudt et al., 1988, Science 241:577-580; Bock et al., 1992, Nature 355:564-566; Tuerk et al., 1992, Proc. Natl. Acad. Sci. USA 89:6988-6992; Ellington et al., 1992, Nature 355:850-852; U.S. Patent No. 5,096,815, U.S. Patent No. 5,223,409, and U.S. Patent No. 5,198,346, all

to Ladner et al.; Rebar and Pabo, 1993, Science 263:671-673; and PCT Publication No. WO 94/18318.

In a specific embodiment, screening can be conducted out by contacting the library members with a Nr-CAM protein (or nucleic acid or derivative) immobilized on a solid phase and harvesting those library members that bind to the protein (or nucleic acid or derivative). Examples of such screening methods, termed "panning" techniques are described by way of example in Parmley and Smith, 1988, Gene \frac{73}{2}:305-318; Fowlkes et al., 1992, BioTechniques \frac{13}{2}:422-427; PCT Publication No. WO 94/18318; and in references cited hereinabove.

In another embodiment, the two-hybrid system for selecting interacting proteins in yeast (Fields and Song, 1989, Nature 340:245-246; Chien et al., 1991, Proc. Natl.

15 Acad. Sci. USA 88:9578-9582) can be used to identify

molecules that specifically bind to a Nr-CAM protein or derivative.

5.10. ANIMAL MODELS

20 The invention also provides animal models. In one embodiment, animal models for diseases and disorders involving cell hypoproliferation (e.g., as described in Section 5.8.1) are provided. Such an animal can be initially produced by promoting homologous recombination between a Nr-25 CAM gene in its chromosome and an exogenous Nr-CAM gene that has been rendered biologically inactive (preferably by insertion of a heterologous sequence, e.g., an antibiotic resistance gene). In a preferred aspect, this homologous recombination is carried out by transforming embryo-derived stem (ES) cells with a vector containing the insertionally inactivated Nr-CAM gene, such that homologous recombination occurs, followed by injecting the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, followed

by the birth of the chimeric animal ("knockout animal") in which a Nr-CAM gene has been inactivated (see Capecchi, 1989, Science 244:1288-1292). The chimeric animal can be bred to produce additional knockout animals. Such animals can be mice, hamsters, sheep, pigs, cattle, etc., and are preferably non-human mammals. In a specific embodiment, a knockout mouse is produced.

Such knockout animals are expected to develop or be predisposed to developing diseases or disorders involving cell hypoproliferation. Such animals can be used to screen for or test molecules for the ability to promote proliferation and thus treat or prevent such diseases and disorders.

In a different embodiment of the invention, transgenic animals that have incorporated and express a functional Nr-CAM gene have use as animal models of diseases and disorders involving cell hyperproliferation or malignancy. Such animals are expected to develop or be predisposed to developing diseases or disorders involving cell hyperproliferation (e.g., malignancy) and thus can have use as animal models of such diseases and disorders, e.g., to screen for or test molecules (e.g., potential anti-cancer therapeutics) for the ability to inhibit overproliferation (e.g., tumor formation) and thus treat or prevent such diseases or disorders.

25 The following examples are provided for the purposes of illustration only and are intended to limit the scope of the invention in any manner.

6. EXAMPLE: ISOLATION OF THE Nr-CAM GENE FROM AND CHARACTERIZATION OF ITS EXPRESSION IN HUMAN GLIOBLASTOMA MULTIFORME TUMOR TISSUE

In this study, the role of Nr-CAM, in brain 5 tumorigenesis was characterized.

6.1. MATERIALS AND METHODS

6.1.1. HUMAN TISSUES AND CELL LINES

Tissue samples of brain and non-brain tumors were
procured from the tissue bank maintained by Pacific Northwest
Cancer Foundation, Northwest Hospital, and from resources at
the Mayo Clinic (Rochester, MN). Brain tumor cell lines
astrocytoma grade IV (CCF-STTG1), astrocytoma grade III (SW
1738), neuroblastoma (IMR-32), medulloblastoma (D283 Med),
glioma (Hs 683), neuroectodermal (PFSK-1), GM(DBTRG-05MG)
were purchased from the ATCC (Rockville, MD). Fetal normal
human astrocytes (FNHAs) were purchased from Clonetics (San
Diego, CA). All cell lines were cultured under the
conditions recommended by the ATCC or Clonetics.

20 6.1.2. DIFFERENTIAL DISPLAY POLYMERASE CHAIN REACTION (DD-PCR)

In order to isolate and clone genes differentially expressed in normal brain tissue (NBT) and glioblastoma multiforme tissue (GMT), the technique of Different Display-PCR (DD-PCR) was utilized. (Examples of protocols of DD-PCR may be found in Sehgal et al., 1997, J. Surg. Oncol. 64:102-108; Sehgal et al., 1997, J. Surg. Oncol. 65:249-257; Sehgal et al., 1997, Int. J. Cancer 71:565-572 (Sehgal, 1997b); Sehgal et al., 1996, Exp. Lung. Res. 22:419-434).

NBT and GMT were obtained from the same region of the brain. Total RNA was isolated and first strand cDNA synthesis was carried out using the first strand cDNA synthesis kit from Clontech (Palo Alto, CA) using BT3-2

primer (5'T [T] 18NG3'). Approximately 125 ng of first strand cDNA synthesis product were used for carrying out PCR. DD-PCR was carried out using (λ P³²) end-labeled BT3-2 primer and BT10 (5-NGCTGCTCTCA TACT-3') primer using cDNA from NBT or GMT tissue in duplicate under the conditions described previously (Sehgal et al., 1997a). PCR products were run on a 6% sequencing gel. Bands that showed differential expressions were cut out, and DNA was eluted and cloned into a PCRII vector Invitrogen, San Diego, CA). Positive clones were screened by PCR and sequenced using the Sequenase version 2.0 sequencing kit from Amersham/USB (Cleveland, OH).

6.1.3. GENE SPECIFIC RT-PCR

To confirm differential expression of clones isolated by DD-PCR, gene-specific RT-PCR technique was 15 carried out as described previously (Sehgal et al., 1997b). hNr-CAM specific primers (5'-AACATATGGGTAGAGAGTATATTT-3' (SEQ. ID No. 9); and 5'-CTTTGCATTCCAGTTCATATTAA-3' (SEQ. ID. NO. 10) were used for PCR. This PCR results in a 250 bp product at the 3' end of the hNr-CAM gene. For EGFR, gene-specific 20 primers (5'-TGTGGTGACAGATCACGGCT-3' (SEQ. ID. No. 11) and 5'-CAGCTCAAACCTGTGATTTCC-3') (SEQ. ID. No. 12) were used for PCR and an internal primer (5'-AATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGC TACGAATATTAAACACTT-3') (SEQ. ID. No. 13) for Southern blot analysis. As a control for PCR, D1-2 (mitochondrial Cytochrome C oxidase subunit 1 gene, Accession Number D38112), a housekeeping gene, which is expressed in both NBT and GMT, was used. PCR was carried out using D1-2-specific primers (5'-CGGAGCAATATGAAATGATCT-3' (SEQ. ID. NO.: 14) and

5'-GCAAATACAGCTCCTATTG-3') (SEQ. ID. NO.: 15), resulting in a

200 bp product. PCR for all 3 genes was carried out using
Taq DNA polymerase under the conditions recommended by Qiagen
(Chatsworth, CA). PCR product was then run on a 2% agarose

gel and transferred onto a Hybond N+ nylon membrane using standard Southern blotting conditions, as described previously (Sehgal *et al.*, 1997a). Hybridization was done at 42°C using hNr-CAM, EGFR and D1-2-specific probes. EGFR,

- 5 hNr-CAM and D1-2-specific probes were prepared by multiprime labeling (Amersham, Arlington Heights, IL) of hNr-CAM-specific primers (5'-
 - GCTGTATGTTAGTATTATGAGAATAGTTACAGCAAAAACATAA CTCAGT-3') (SEQ. ID. No.: 16) or D1-2-specific primer (5'-
- 10
 17). These primers are internal to the primers used for PCR, and they do not carry any of the primer sequences used in the PCR. Primer sequences were checked for homologous sequences using the DNA BLAST program of NCBI (National center for Biotechnology Information, Bethesda, MD) prior to usage. Quantitation of the signal on Southern blot was carried out using the ImageQuaNT program of Molecular Dynamics (Sunnyvale, CA). This protocol was used to quantitate expression of EGFR, D4-1 or D1-2 in brain tumor
- 20 demonstrated previously that this gene-specific RT-PCR technique is semi-quantitative (Sehgal et al., 1997a, b).

cell lines, FNHA and selected tumor tissues. We have

6.1.4. NORTHERN BLOT ANALYSIS

Multiple Normal Human tissue blots (MNHTB) were purchased from Clontech (Palo Alto, CA). These blots contained 2μg of pure polyA+ mRNA. MNHTBs were prehybridized in express hybridization buffer solution (Clontech) for 3-4 hours. Hybridization was done with multiprime labeled 179 bp D4-1 probe. After autoradiographic exposure, the probe was washed from the blot and then hybridized with human β actin probe (Clontech). Quantification of expression of hNr-CAM and β actin was done using the ImageQuaNT program.

Expression of Nr-CAM in different regions of normal brain and cell lines formed from tumor tissues was assessed.

6.1.5. QUANTITATION OF NORTHERN AND SOUTHERN BLOTS

year performed using the ImageQuaNT volume quantitation program. Volume quantitation calculates the volume under the surface created by a 3-D plot of pixel locations and pixel values. We quantitated the volume (the integrated intensity of all pixels in the spot excluding background) of D1-2 bands in Northern or Southern blots. These pixel values are then normalized with pixel values in the bands of housekeeping genes (D1-2 or β actin) and are referred to as "relative expression" in the figures. The subjective terms "low", "medium" and "high" refer to relative expression and are based on hNr-CAM expression in normal brain as "low" and in tumor brain as "high".

6.1.6. IN SITU HYBRIDIZATION

The technique of *In situ* hybridization was done as 20 described previously (Wilkinson, 1992 In Situ Hybridization, A practical approach. NY: Oxford University Press).

Briefly, 6μm formalin fixed, paraffin embedded human brain tumor sections were deparaffinized by 2 washes in xylene, followed by rehydration through graded concentrations of ethanol from 100% to 70%. These were then washed in PBS and treated with Proteinase K (25 mg/ml for 10 minutes), followed by fixation in 4% paraformaldehyde. After incubation in 0.25% acetic anhydride/0.1 M TEA (Tri-Ethyl Acetic acid), sections were dehydrated through graded concentrations of ethanol from 70% to 100% and prehybridized for 2 hours at 55°C in 50% formamide, 5xSSC pH 4.5, 50μg/ml tRNA, 50 μg/ml heparin, and 1% SDS. Sections were hybridized with 1 μg/ml

DIG-(Digoxygenin) labeled antisense or sense probes for 18 hours at 55°C.

Probes, sense and anti-sense, were synthesized with the Genius 4 kit (Boehringer Mannheim, Indianapolis, IN) 5 using the T3 and T7 promoters of a PCR template derived from human Bravo/Nr-CAM sequences corresponding to bases 3731-3754 and 4101-4114. See Figure 2A in which BT180 represents the 5' primer for the probe corresponding to nucleotides 3731-3754 and BT181, the 3' primer for the probe corresponding to nucleotides 4104-4114. Following hybridization, slides were 10 washed in 50% formamide, 2xSSC pH 4.5, 1 % SDS at 50°C, treated with 5 μ g/ml RNase A for 30 minutes at 37°C, and washed in 50% formamide, 2xSSC pH 4.5 at 50°C. Sections were pre-blocked in 10% normal sheep serum (Sigma, St. Louis, MO) and incubated with a 1:2000 dilution of alkaline phosphate 15 conjugated anti-dioxigenin Fab fragments (Boehringer Mannheim) 18 hours at 4°C. For detection, slides were incubated with NBT/BCIP (5-Bromo-4-chloro-3-indilylphosphate, 4-toluidine salt) in the dark for 46 hours. counter staining with eosin Y, slides were mounted with 20 Permount and visualized using an Axioskop (Carl Zeiss, Thornwood, NY) routine microscope.

6.1.7. GENOMIC SOUTHERN BLOT

NIH3T3, astrocytoma III, glioma and glioblastoma

cells were grown in 100 mm diameter plates until 90%

confluent under the conditions recommended by the ATCC.

Genomic DNA was isolated using a DNA isolation kit from

Puragen (Research Triangle Park, NC); 10 μg of DNA were cut

with 50 U of EcoRI restriction enzyme from GIBCO BRL

(Gaithersburg, MD) and run on a 1% agarose gel. DNA was

transferred onto a Hybond N+ nylon membrane (Amersham) using

the protocol recommended by Puragen. After incubating the

membrane in pre-hybridization solution, hybridization was

conducted using a 375 bp hNr-CAM probe (nucleotide position 3731-4126) (Lane et al., 1996, Genomics 35:456-465) labeled (1 X 104 CPM/ml) using the multiprime labeling kit from Amersham. Membrane was washed in 0.1 X SSC/0.1% SDS at 24°C for 30 mins and then exposed to Kodak X-ray film.

6.2. RESULTS

6.2.1. ISOLATION OF HUMAN Nr-CAM FROM AND DIFFERENTIAL EXPRESSION OF Nr-CAM IN GLIOBLASTOMA

Using the modified technique of DD-PCR as described in Section 6.1.2., a cDNA fragment, designated D4-1, was identified which is over-expressed in glioblastoma multiform tissue (GMT) as compared with normal brain tissue (NBT). Figure 1 demonstrates that D4-1 is over-expressed in GMT.

15 The band designated C in Figure 1 is the cDNA of the D1-2 gene, a control gene present in both GMT and NBT.

The D4-1 band was isolated from the gel and cloned into the pCRII vector from Invitrogen as described in Section 6.1.2. Sequence identity analysis indicated that this cDNA is identical to the last 38 bases at the 3' end of the previously isolated hNr-CAM gene. Figure 2C presents the results of sequence identity analysis together with a comparison of the sequence of the isolated hNr-CAM with that of the rat Nr-CAM. The isolated D4-1 cDNA has 99.2% sequence identity (homology) to the rat Nr-CAM. Thus, the sequence identity analysis demonstrates that clone D4-2 is hNr-CAM.

In situ hybridization conducted as described in Section 6.1.6 using an anti-sense Nr-CAM confirmed differential expression of hNr-CAM in glioblastoma as compared to normal brain tissue. Results are presented in Figures 3(a-f).

As shown in Figures 3a and d, strong expression of hNr-CAM is observed in a number of cells from 2 GMT samples,

using the anti-sense hNr-CAM probe. Low or no signal was observed when hNr-CAM sense probe was used on serial sections (Fig. 3b,e). The NBTs did not show any signal with the hNr-CAM anti-sense probe (Fig. 3c,f). This experiment indicates that hNr-CAM is differentially over-expressed in GMT as compared to NBT.

6.2.2. EXPRESSION OF Nr-CAM IN HUMAN TUMOR TISSUES

As shown in Figures 4 (A and B), hNr-CAM is

expressed at high levels in glioblastoma IV and glioma tissue
as compared to NBT using RT-PCR. Low or no expression of
hNr-CAM was observed in recurrent meningioma, meningioma,
neuroblastoma, recurrent malignant glioma, melanoma, breast
tumor, benign prostate tissue or NBT. In one of the GMT
samples studied, no hNr-CAM expression was observed. High
expression of hNr-CAM was observed in NBT undergoing
extensive gliosis. This is an unusual observation, and the
reason for it is not known at present. D1-2 is a
housekeeping gene that has been used previously as an
internal control in RT-PCR (Sehgal et al., 1997a,b).

The expression of hNr-CAM in NBT and astrocytoma tumor tissue was evaluated using RT-PCR. As shown in Figures 5 (A and B), a high level of hNr-CAM is expressed in astrocytoma as compared to NBT. The expression pattern of EGFR, a known brain tumor marker, was investigated from this set of cDNA. As shown in the middle and bottom panels of Figures 5A and in 5B, expression of hNr-CAM is higher in tumor than is EGFR.

6.2.3. EXPRESSION OF Nr-CAM IN BRAIN TUMOR CELL LINES

Expression of hNr-CAM in cell lines derived from several different kinds of human brain tumor was investigated. As shown in Figures 6 (A and B), Nr-CAM is expressed at high levels in astrocytoma IV, glioma,

30

glioblastoma and neuroectodermal tumor cell lines as compared to NBT. Low or no expression of hNr-CAM was observed in cell lines derived from astrocytoma III, medulloblastoma, neuroblastoma and NBT. The hNr-CAM transcript was not detected in FNHAs or NIH3T3 cells (data not shown).

6.2.4. EXPRESSION OF Nr-CAM IN NORMAL HUMAN BRAIN

cord were studied for hNr-CAM expression. As shown in

Figures 7 (A and B), a natural transcript of hNr-CAM (7.5 kb)

was observed in cerebellum, occipital lobe, cerebral cortex and frontal lobe at a higher level as compared to spinal cord, medulla, temporal lobe or putamen. Expression of hNr-CAM detected is low as compared to tumor cell lines (Figures 8 (A and B) or the other tumor-associated gene (D2-2)

previously studied. Expression of hNr-CAM is highest in kidney, and no significant difference in expression between fetal and adult tissues was observed (data not shown).

6.2.5. EXPRESSION OF Nr-CAM IN HUMAN TUMOR CELL LINES

As shown in Figures 8 (A and)B, hNr-CAM was expressed at high levels in melanoma G361, lymphoblastic leukemia (MOLT-4) and Burkitt's lymphoma Raji cell lines. A low level of hNr-CAM expression was observed in promyelocytic leukemia (HL-60), HeLa cell S3, chronic myelogenous leukemia (K-562), colorectal adrenocarcinoma (SW480) and lung carcinoma (A549). All of the cell lines studied herein expressed hNr-CAM mRNAs that are 1.4 kb as compared to the 7.5 kb transcript expressed in normal brain (Figures 5 (A and B). HeLa cells S3 express low levels of both transcripts.

Melanoma G361 express high levels of the 7.5 kb and low levels of the 1.4 kb transcript, suggesting alternative splicing of hNr-CAM mRNA during tumorigenesis.

Using Northern blot analysis as described in Section 6.1.4, expression of Nr-CAM was investigated in 8 different human tumor cell lines.

6.2.6. EXPRESSION OF Nr-CAM IN HUMAN BRAIN TUMORS USING IN SITU HYBRIDIZATION

In situ hybridization as described above was performed on a panel of 20 different brain tumors using antisense and sense hNr-CAM-specific probes. Results from this study are shown in Table 2.

10

5

15

20

25

30

TABLE 2

EXPRESSION OF HNR-CAM IN HUMAN BRAIN TUMORS: PRESENCE
OR ABSENCE OF HNR-CAM IS SHOWN BY POSITIVE OR NEGATIVE SIGNS

_	TISSUE	hNr-CAM
5	Glioblastoma multiforme	+
		+ '
		÷
	-	+
		-
		-
		-
	·	-
	Genicystic anaplastic astrocytoma	-
	Meningotheliomatous meningioma	+
	Normal brain	-
	·	-
		-
		-
	Glioma III/IV	+
	Pilocystic astrocytoma	+
	Malignant glioma	-
	Lipidized meningioma	+
	Meningioma syncytial type	+
	Meningioma	-
5.	Fibroblastic meningioma	-
	Meningotheliomatous meningioma, grade I	+
	Genicystic astrocytoma	-
	Oligodendroglioma, grade II	+

30

Eleven of the 20 tumors (55%) showed positive signal for hNr-CAM. Four NBT samples did not show any hNr-

CAM expression. None of the early grade astrocytomas and only 50% of the highly anaplastic astrocytomas showed Nr-CAM expression. Similar hNr-CAM expression was observed in brain tumor cell lines (Figures 6 (A and B)). hNr-CAM expression was observed in 50% of meningiomas tested and one oligodendroglioma. These results demonstrate that expression of hNr-CAM is prevalent in malignant glioma tissue.

6.2.7. GENE AMPLIFICATION

Genomic Southern blot was performed as described in Section 6.1.7 on 3 brain tumor cell lines (astrocytoma III, glioma and glioblastoma) and the NIH3T3 cell line. As shown in Figures 9 (A and B), no change in the genetic level of hNr-CAM was observed in the 4 cell lines tested.

These results indicate that the over-expression of hNr-CAM in brain tumors is not due to gene amplification.

A 1.4 kb transcript of hNr-CAM was observed presently in tumor cell lines as compared to a 7.5 kb transcript in normal brain (Lane et al., 1996). This may indicate that the 7.5 kb transcript for hNr-CAM generates a 20 1.4 kb transcript that could translate into a small version of the hNr-CAM protein, which may be tumor-specific. On the basis of data presented in this study, we conclude that hNr-CAM is over-expressed in human malignant brain tumors and that it is useful to serve as a marker for detection and for therapy.

7. EXAMPLE: EFFECT OF REGULATING Nr-CAM EXPRESSION IN GLIOBLASTOMA

In order to assess the functional role of Nr-CAM in brain tumorigenesis, the effects of over-expressing Nr-CAM in the anti-sense direction were examined in a glioblastoma cell line.

7.1. MATERIALS AND METHODS

7.1.1. CLONING OF ANTISENSE Nr-CAM

The full length clone for Nr-CAM was provided by William Dryer (CalTech). Three different portions of the 5 hNr-CAM gene were cloned in the antisense direction.

To obtain antisense "Nr-CAM 1/3 clone", Nr-CAM 1/3 (corresponding to nucleotides beginning at nucleotide 119 and ending at nucleotide 1434 of Figure 2A) was amplified using primers BT306 (5' TAGATACAACTAGTCTAATGCAGCTTAAAATAATGCC 3') (SEQ.ID. No.: 18) and BT307 (5'

- AGATAGATCCGCGGATATCCATATTCATTAGAGGCATTG 3') (SEQ. ID. No. 19)

 (see Figure 2A) and cloned into precut pCMVneo vector cut with SACII and SPEI restriction enzymes. PCR amplification was carried out for 1 cycle at 94°C 3 min, 61°C 1 min, 72°C 4 min, then for 30 cycles at 94°C 1 min, 61°C 1 min, 72°C 4 min followed by 1 cycle at 94°C 1 min, 61°C 1 min, 72°C 10 min. The PCR product was cut with Spel and Sacil, and cloned in the antisense direction into the pCMV-neo vector precut with SacII and Spel enzymes. Orientation of the hNr-CAM gene was confirmed by restriction digestion of specific enzymes.
- A 1.3Kb fragment of the hNr-CAM gene (spanning the first 1/3 part of the gene) was cloned in the antisense direction into precut pCMVneo vector (See Figure 2D). This vector contains a constitutively active cytomegalovirus promoter and has been used in the past to over-express genes in cells (Huang, et al., 1997, Int J Cancer, 72:102-109). This clone was also termed "pCMV-1/3Nr-AS". The pCMV-neo or pCMV-1/3Nr-AS were then transfected into the 5GB glioblastoma cell line (ATCC# 2020-CRL), and selected in G418.

20 This clone was termed "pCMV-1/3Nr-AS".

To obtain anti-sense Nr-CAM 2/3 clone, Nr-CAM 2/3

(corresponding to nucleotides 1410-2746 of Figure 2A) was amplified using primers BT308 (5'

TAGATACAACTAGTCAATGCCTCTAATGAATATGGATA 3') SEQ. ID. No.: 20);

and BT309 (5' AGATAGATCCGCGGAATAGTAAATCCGATAGCCTTGTA 3') (SEQ. ID. No.: 21) cut with Spe I and SacII, and cloned into precut pCMVneo vector cut with Sac II and Spe I enzymes.

7.1.2. TRANSFECTION OF GLIOBLASTOMA (GB)

5

GB cells were plated at an approximate density of 3 x 104. 24 hours after plating the cells were washed with serum-free media and transfected with lipofectamine reagent plus plasmid DNA diluted in 1 ml total of serum-free media. Cells were incubated at 37°C for hours after which the 10 reagent was replaced with media containing 10% FBS. Cells were incubated at 37°C for 72 hours. Media was changed adding 1000 μ g/ml G418 a selective media to select for resistance to neomycynin (GIBCO-BRL). Cells were incubated at 37°C for 72 hours. Media was changed adding 1000 μ g/ml 15 G418. Cells were incubated at 37°C for 96 hours. changed adding 1000 µg/ml G418. Cells were incubated at 37°C for 72 hours. At this point all the cells in the control plate were dead. At this point, media was changed, adding 400 μ g/ml G418. Cultures were maintained at 400 μ g/ml G418 20 indefinitely, changing media every 72-96 hours.

7.1.3. CELL MORPHOLOGY

During maintenance, cells transfected as described in Section 7.1.2 above, were split at a ratio of 1:2-1:3.

After 72-96 hours, photographs of the cells were taken to compare the morphology.

7.1.4. GROWTH ASSAY

On Day 0, 5GB, 5GBpCMV neo, 5GBNr-CAM 1/3, 5GBNr-CAM 2/3 cells were trypsinized, counted on a Coulter Counter and plated at a density of 1 x 10⁴ cells/60 mm dish. 12 plates of each condition were plated. At each time point 3

plates of each condition were counted on the Coulter Counter and the counts averaged.

7.1.5. SOFT AGAR ASSAY

Soft agar assay, a common in vitro phenotype of transformation, was performed as described previously (Huang et al., 1995, Cancer Research 55:5054-5062). Briefly, 5GB cells that were transfected with vector alone and with Nr-CAM in anti-sense direction were trypsinized. Approximately, 1X10⁵ cell were mixed with 0.26% agar. Cells were then plated on top of a layer of 0.65% agar in 60mm petri dishes and incubated 37°C for 2 weeks. Cells were fed with one ml media containing 10% FBS after 1 week. Colonies were stained and counted under the inverted light microscope.

7.1.6. NORTHERN BLOT ANALYSIS

PolyA+ mRNA was prepared from 5GB glioblastoma cells transfected with pCMVneo or pCMV-1/3Nr-AS cells using the Quick Prep mRNA Purification Kit (Pharmacia Biotech, Piscataway NJ). $2\mu g$ of polyA+ RNA was run in two lanes.

- 20 Approximately 1x10° cpm/ml of labeled hNr-CAM probe was added to the blot and hybridized for 18 hours at 68°C. After washing in 0.1% SSC, 0.1% SDS, the blot was visualized using a Phosphor Imager (Molecular Dynamics) and the signal quantified using ImageQuant software (Molecular Dynamics).
- 25 After quantification, the blot was stripped by incubating at 90°C for 4 minutes in 0.1% SDS solution. The membrane was then prehybridized for 6 hours at 68°C and then treated with a probe for β-actin at a concentration of 5x10⁵ cpm/ml for 18 hours at 68°C. After washing, the blot was analyzed using the Phosphor Imager as described above. The volumes of the images for each probe were compared and normalized for β-

actin signal.

7.2. RESULTS

7.2.1. EFFECT OF EXPRESSION OF ANTI-SENSE Nr-CAM ON MORPHOLOGY

To study the role of the Nr-CAM gene in cell transformation, Nr-CAM was over-expressed in 5GB cells in the anti-sense direction. The Nr-CAM antisense construct designated "Nr-1/3AS" was used. Approximately 10µg of pure DNA was transfected onto two 60mm-diameter petri-dishes containing 10,000 cells using lipofectamine (Gibco/BRL). Transfected cells were selected in G418 (1000 μ g/ml) for 2 10 weeks. After 3 weeks, cells were maintained in $400\mu\text{g/ml}$ G418. Cell morphology was observed under the inverted light microscope and cell proliferation properties of transfected cell were analyzed by counting cell number at various intervals. Results are presented in Figures 10A and B. 15 Figure 10A shows 5GB pCMVneo cells 96 hours after media change. Figure 10B shows 5GBNr-CAM 1/3 cells 96 hours after media change. No change in cell growth and morphology was observed in glioblastoma cells transfected with pCMV-neo vector (control) but cells transfected with Nr-CAM 1/3 in anti-sense direction (pCMV-neoCA) showed a change in cell morphology and slower cell proliferation (see Figure 10B). Our results indicate that Nr-CAM over-expression in the antisense direction blocked Nr-CAM gene further in the 5GB glioblastoma cell line. This result strongly suggests that Nr-CAM expression is required for continuous proliferation of 25 5GB cells.

In another set of experiments, the expression of antisense expression of Nr-CAM on GB morphology was evaluated in 5GB cells using the pCMV-1/3 Nr-AS construct.

5GB glioblastoma cells were plated at an

30 approximate density of 3x10⁴ • 24 hours after plating the

cells were washed with serum-free media and transfected withlipofectamine reagent plus plasmid DNA diluted in 1 ml total

of serum-free media. Cells were incubated at 37°C for 5 hours after which the reagent was replaced with media containing 10% FBS. Cells were incubated at 37°C for 72 hours. Media was changed to the one containing $1000~\mu\text{g/ml}$ 5 G418. Cells were incubated at 37°C for 7 days. At this point, media was changed to the one containing $200~\mu\text{g/ml}$ G418. Cultures were maintained at $200~\mu\text{g/ml}$ G418 indefinitely, changing media every 72-96 hours. 96 hours after plating cells, photographs were taken to compare the morphology and are presented in Figures 15 (A-D).

After transfection of antisense hNr-CAM (pCMV-1/3Nr-AS), glioblastoma cells were selected in G418 media for two weeks (1000μg/ml). Untransfected 5GB cells (PCMV-neo) were used as controls. Cell morphology was compared between pCMV-neo or pCMV-1/3Nr-AS transfected cells after four weeks of selection. The glioblastoma cells transfected with antisense hNr-CAM became spindle shaped and showed neurite outgrowth (compare Figures 15 A and B with Figures 15 C and D). 5GB cells transfected with pCMV-1/3Nr-AS were grown in culture for 3 weeks and they demonstrated lack of density dependent inhibition of cell proliferation.

One unique observation in maintaining the hNr-CAM antisense transfected cells is that when fresh media containing 10% fetal bovine serum (FBS) is added to these cells, their morphology changes temporarily to one that is similar to pCMV-neo transfected 5GB cells. Both pCMV-neo and pCMV-1/3Nr-AS transfected cells were treated with different concentrations of serum (0.1, 1, 2 and 5% FBS). As shown in Figure 16, 2% FBS is sufficient to cause a change in pCMV-1/3Nr-AS transfected 5GB cells similar to pCMV-neo transfected cells. No change in cell morphology was observed in cells transfected with pCMV-neo. This result suggests that one or a combination of more than one growth factors in the

serum transiently reverses the morphology enforced by the anti-sense *hNr-CAM*.

7.2.2. EFFECT OF EXPRESSION OF ANTI-SENSE Nr-CAM ON CELL PROLIFERATION

The effect of Nr-CAM expression in the anti-sense direction on glioblastoma proliferation was evaluated as described above in Section 7.1 using anti-sense Nr-CAM 1/3 or pCMV-1/3Nr-AS. Results are illustrated in Figure 11 and Figure 17.

As shown in Figure 11, expression of anti-sense Nr-CAM inhibited proliferation of GB cells compared to GB cells containing vector only (GB/pFCS).

As shown in Figure 17, 5GB cells transfected with pCMV-1/3Nr-AS proliferate slowly as compared to pCMV-neo
15 transfected cells. This result clearly demonstrates that hNr-CAM is required for continuous proliferation of cells.

Even though 5GB cells transfected with pCMV-1/3Nr-AS proliferate slowly, they maintained their spindle shape morphology.

20

5

7.2.3. SOFT AGAR COLONY FORMATION OF Nr-CAM ANTISENSE EXPRESSING CELLS

Results presented in Figure 12 demonstrate that expression of Nr-CAM in the anti-sense direction (GM-Anti-Nr-CAM) inhibits the number of soft agar colonies compared to results observed with non-transfected GB cells (GB) and GB cells transfected with control plasmid only (GB-PFS). As illustrated, overexpression of antisense hNr-CAM caused 81% inhibition in number of soft agar colonies formed. Colonies formed by untransfected 5GB cells and control transfected 5GB cells with pCMV-neo were larger than those expressing hNr-CAM antisense.

7.2.4. EXPRESSION OF Nr-CAM IN hNr-CAM-ANTISENSE EXPRESSING CELLS

The expression of hNr-CAM in cells expressing hNr-CAM antisense was evaluated using the Northern blot analysis technique described in Section 7.1.6.

As shown in Figure 13, over-expression of the hNr-CAM anti-sense caused approximately 60% reduction in the native hNr-CAM expression. A logical explanation of this could be that in antisense hNr-CAM transfected cells, the natural transcript is made constitutively and a percentage of it is detected by Northern blot analysis regardless of RNase mediated degradation of antisense hNr-CAM bound to natural transcript (See Figure 14).

7.2.5. CELL CYCLE ANALYSIS OF CELLS EXPRESSING hnr-CAM ANTISENSE

To study cell cycle status of 2CMV-neo and pCMV1/3Nr-AS transfected cells, flow cytometry was performed on
5GB cell cycle status. Approximately 6x10⁶ 5GB cells
transfected with pCMVneo, or pCMV-Nr1/3AS cells were
harvested by trypsinization followed by fixation in 80%
ethanol (vol/vol) fixative (Sigma, St. Louis MO) and
incubated for 24 hours at -20°C. The cells were then stained
with Propidium Iodide for 30 minutes at room temperature in
the dark. After filtering to remove debris, the cells were
read on a FACS Calibur cell sorter (Becton Dickinson).

Twenty thousand gated events were counted and the results
analyzed using ModFit Lt 2.0 software (Becton Dickinson).

Results are presented in Table 3.

30

15

TABLE 3

EFFECT OF HNR-CAM ANTISENSE OVER-EXPRESSION

	CELL CYCLE PHASES	% OF CELLS
5	pCMV-neo Transfected Cells	
	G0-G1	72.76
	G2 - M	10.00
	s	17.24
10	. G2-G1	1.83
	pCMV-1/3Nr-AS Transfected Cells	
	G0-G1	89.98
	G2-M	3.66
	S	6.36
15	G2-G1	1.83

As shown in Table 3, a 63% decrease in S phase, 20% increase in G0-G1 phase and a 63.4% decrease in G2-M was observed in pCMV-1/3Nr-AS transfected cells as compared to pCMV-neo transfected cells. This set of results clearly demonstrates that antisense hNr-CAM causes a lengthening of specific phases of 5GB glioblastoma cells, i.e., causes lengthening of the cell cycle.

7.2.6. EFFECT OF ANTISENSE hnr-cam on MIGRATION AND INVASION

Glioblastoma cells are highly invasive and they penetrate into surrounding normal brain tissue during their genesis (Kleihues and Cavenee, Pathology and genetics of tumors of the nervous system. Lyon, France: International Agency for Research on Cancer, 1997). To determine if antisense hNr-CAM could alter the migration capacity of glioblastoma cells, a cell migration assay was performed on

25

5GB cells transfected with pCMV-neo or pCMV-1/3Nr-AS. Equal number of cells (1x10⁶) were plated on a 8μm pore size polycarbonate membrane filter. Cells that migrated through the membrane after 3 days were counted after fixation and staining with hematoxylin. Results are shown in Figure 18.

As shown in Figure 18, anti-sense hNr-CAM over-expression caused a 30% inhibition in the migration ability of 5GB glioblastoma cells.

In addition, an invasion assay was performed (See, Ridder and Calliauw, 1992, Neurosurgery, 31:1043-1048) to determine if antisense hNr-CAM could inhibit the invasion properties of glioblastoma cells.

Briefly, 825ng of ECM gel was coated on to $8\mu m$ pore size polycarbonate membrane filter. Equal number of 5GB cells $(1x10^4)$ were plated on to the ECM gel. Cells that migrated through the ECM gel after 4 and 7 days were counted after fixation and staining with hematoxylin. Results are presented in Figure 19.

As shown in Figure 19, approximately 90% inhibition of cell invasion was observed in pCMV-1/3Nr-AS transfected 20 5GB cells as compared to pCMV-neo transfected cells.

7.2.7. EFFECT OF RADIATION OF CELLS ANTISENSE hnr-CAM

radiation. To determine if treatment antisense hNr-CAM
transfected glioblastoma cells are UV radiation sensitive,
the following experiment was conducted. Approximately 1x10⁴
glioblastoma cell (transfected either with pCMV-neo or pCMV1/3Nr-AS) were plated in triplicates in 60mm diameter
petridishes. Cells were then exposed to 100 units of UV
radiation. As observed 65% of antisense hNr-CAM transfected
cells died as compared to 27% death in pCMV-neo transfected
cells.

The percentage of surviving cells undergoing apoptosis was also determined. To do so, the UV radiation experiment was repeated and cells undergoing apoptosis were identified using Apoptosis kit from BMB (Indianapolis, IA). Results are presented in Figure 20.

As shown in Figure 20, a 17 fold increase in the number of cells undergoing apoptosis was observed. These results clearly suggest that antisense hNr-CAM over-expression caused 5GB glioblastoma cells to become more sensitive to UV radiation.

10

20

25

7.2.8. ANTISENSE bnr-CAM INHIBITS GB TUMOR GROWTH IN VIVO

In one experiment, designated "Experiment 1", 5 nude mice were subcutaneously with 3.0x10⁶ 5GB glioblastoma 15 cells (pCMV-neo or pCMV-1/3Nr-AS transfected). Results are presented in Table 4.

As shown in Table 4, tumor growth was observed in three of the five mice that were injected with 5GB (pCMV-neo transfected) cells. No tumor growth was observed in mice injected with pCMV-1/3Nr-AS transfected 5GB cells.

To increase the efficiency of this type of experiment, in another experiment, designated "Experiment 2", 1x10⁷ 5GB cells (transfected with pCMV-neo or pCMV1/3Nr-AS) were injected into seven nude mice. Tumor size was measured 38 days post-injection. Results are presented in Table 4.

In Experiment 2, six of seven mice injected with cells transfected with pCMV-neo developed tumor. No tumor growth was observed in mice that were injected with cells transfected with pCMV-1/3Nr-AS vector.

Photographic illustration of three examples (each)

30 of mice injected with pCMV-neo or pCMV-1/3Nr-AS vectors is

presented in Figure 21.

Results from this experiment clearly demonstrated that anti-sense hNr-CAM inhibits tumorigenic properties of 5GB glioblastoma cells in vivo. Injection of antisense hNr-CAM expressing glioblastoma cells caused inhibition of tumor formation.

TABLE 4

THE EFFECT OF ANTISENSE NNR-CAM

EXPRESSION ON TUMOR FORMATION IN VIVO

	Experiment 1		Experiment 2	
	Mice injected (5GB glioblastoma cells)		Mice injected (5GB glioblastoma cells)	
15	pCMV-neo	pCMV1/3Nr-As	pCMV-neo	pCMV1/3Nr-AS
	Tumor Volume	(mm³)	Tumor Volume	(mm³)
	400	NT	75	NT ·
	726	NT	650	NT
	936	NT	63	NT
20	NT	NT	196	NT
	NT	NT	365	NT
			196	NT
1			NT	NT .

NT = No Tumor

10

7.2.9. INTRATUMORAL INOCULATION OF PLASMID-EXPRESSING ANTISENSE *bNR-CAM* CAUSED REDUCTION IN GLIOBLASTOMA TUMOR GROWTH

To determine if antisense hNr-CAM could cause reduction in tumor growth in vivo, the effect of direct intra-tumoral injection of an antisense hNr-CAM expressing plasmid mixed with liposomes was analyzed.

^{25 2 =} In experiment 1, tumor size was measured 70 days post injection. In experiment 2, tumor size was measured 38 days post injection.

In a first set of experiments, three athymic nude mice were injected with 1 x 10⁵ 5GB (glioblastoma) cells subcutaneously. 72 days post-implantation, 50 µg of either pCMVneo (control) (one animal) or pCMV1/3Nr-AS (two animals) plasmids were mixed with DMRIE (liopsomes) reagent (Gibco/BRL) and injected twice a week for four weeks around the tumor site. Tumor size was measured twice a week with a caliper and tumor volume was determined. Results are shown in Figure 22A.

As demonstrated in Figure 22A, animals which received anti-sense hNr-CAM injected directly into tmor showed not only slower tumor growth but also tumor regression compared to the control animal.

In another set of experiments, 20 athymic nude mice were implanted with 3x3mm pieces of glioblastoma tumor. 28

15 days post implantation, 300µg of either pCMVneo or pCMV1/3Nr-AS plasmids were mixed with DMRIE reagent (Gibco/BRL) to a final of 300µl volume and injected twice a week for four weeks around the tumor. Control mice were injected with the same volume of 1xPBS or no treatment. Tumor size was

20 measured twice a week with a caliper and tumor volume was determined. Results are shown in Figure 22B.

As shown in Figure 22B, direct intra-tumoral injection of plasmid expressing the antisense hNr-CAM caused slower tumor growth. Results from this set of experiments demonstrate that targeting of the hNr-CAM gene is an advantageous strategy for treating human glioblastoma tumors.

7.2.10. ROLE OF hnr-CAM IS NOT CONFINED TO ONE CELL LINE

As a model to study the role of hNr-CAM in

30 malignant gliomas, we used the 5GB glioblastoma cell line.

Glioblastoma tumor cells are heterogeneous and tumors
isolated from different patients show different genetic

characteristics. Thus, in order to demonstrate the fact that hNr-CAM is a good genetic target for gene therapy of human glioblastomas we demonstrated the anti-tumorigenic properties of antisense hNr-CAM in different glioblastoma cell lines.

Using the antisense hNr-CAM expressing vector (pCMV-1/3Nr-AS), we have not been able to block nHr-CAM expression in GB 1690 glioblastoma cells. This could possibly be due to cellular factors interfering with the binding of the antisense molecule to the appropriate site on hNr-CAM message.

To overcome this problem, we decided to target a different region of the hNr-CAM gene.

We have PCR amplified 1360 bases of the hNr-CAM gene (position 1410-2746) and cloned it in the antisense direction into precut pCMV-neo vector. This clone was termed as pCMV-2/3Nr-AS (Spanning the 2/3rd region of hNr-CAM gene). Preliminary results are shown in Figures 23 (A-C).

The results obtained have demonstrated that pCMV-2/3Nr-AS transfected GB1690 cells changed in cell morphology and formed fewer numbers of soft agar colonies as compared to pCMV-neo transfected cells (Figure 17).

On the basis of this result, it is concluded that hNr-CAM expression requirement for glioblastoma cell proliferation and tumorigenic properties is not confined to just one cell line.

25

7.2.11. <u>ADDITIONAL STUDIES</u>

It is envisaged that anti-sense hNr-CAM can be over-expressed in other human malignant glioma cell lines, such as 1690-CRL, 16-HTB, 138-HTB and other species cells such as, e.g., two rat glioma cell lines, C6 glioma and 9L gliosarcoma. To demonstrate the fact that antisense human Nr-CAM can bind to rat Nr-CAM and inhibit its function, we performed a nucleotide sequence comparison between the rat

and human Nr-CAM sequence. As shown in Figure 24, 87% sequence similarity was observed between the human and rat Nr-CAM nucleotide sequence. On the basis of this result, we conclude that human Nr-CAM antisense molecule is capable of binding to rat Nr-CAM.

It is further envisaged that antisense phosphorothicate oligonucleotides can be used to inhibit the expression of hNr CAM in glioblastoma cells.

Antisense phosphorothioate oligonucleotides can be delivered effectively to several different regions of the brain using high-flow microinfusion technology. Targeting of the hNrCAM gene using antisense phosphorothioate oligonucleotides will be an effective way of treating human glioblastoma tumors in a clinical setting.

As a non-limiting illustrative example, the following is presented. Briefly, we have designed three phosphorothicate oligonucleotides (ODNs) against the translational initiation site of hNr-CAM (see Table 5 below).

Table 5 schematicly illustrates phosphorothioate oligonucleotides (ODAs) for hNr-CAM gene. ODNs H-1, H-2 and 20 H-3 are designed against hNr-CAM; OL-4,OL-5 and OL-6 are random ODNs that can be used as controls. ATG sequence is indicated in bold. Three random ODNs that can serve as controls (OL-4, OL-5 and OL-6) are available commercially from Oliogos Etc. Inc. (Wilsonville, OR).

25 The effect of ODNs on inhibition of hNr-CAM expression can be evaluated using the methodology described previously (Anfossi, et al., 1989 Proc Natl Acad Sci USA, 86:3379-3383). Briefly, 5GB, HTB-16 and GB1690 cells are plated per well in 96-well plates in media without ODNs.

Twenty-four hours later, the culture is media changed to contain a final concentration of 1mmole/L, 3mmol/L, or

10mmol/L ODNs. Control cultures receive fresh culture media without ODNs. After 4-5 days post-transfection, cell

proliferation is analyzed using a cell proliferation assay kit from Promega (Madison, WI). Expression is analyzed using immunocytochemistry methods described previously (Sehgal, et al., 1998, Int. J. Cancer, 76(4):451-458). These oligonucleotides are be tagged with fluorescent tags to ensure their entry into the cells.

TABLE 5 ILLUSTRATIVE ODN'S FOR hNr-CAM

10				ID	SEQ. <u>. No</u> .
	5 'AGGAGTTAAGATGCTAATGCAGC	TTAAAATA ATG CCGAA	AAAGAAGCGCTTATCTGCGGGC3	hNr-CAM	. 22
	3 ' TCCTCAATTCTACGATTAC5 '	H-1	•		23
	3 'ACGTO	GAATTTTATTACGGCT	'5' H-2		24
			3 'TTCTTCGCGAATAGACG5 '	H-3	25
	5 ' ACTAGAGATACAGATCATAT3 '	OL-4			26
15	5 'CATATACGATCGATCGATGC3 '	OL-5			27
	5 GATAGTGCTGATCGATGCTA3	OL-6			28

It is further envisaged that the expression of hNr-CAM gene can be blocked using a replication defective retroviral system to deliver antisense hNr-CAM. Replication defective retroviral systems have been used in the past to deliver genes to a variety of tumor cell types (see, Kondo, et al., 1998, Cancer Res. 68:962-967; Boviaisis, et al., 1994, Human Gene Ther. 5:183-191).

The current state of retroviral gene transfer

technology stems from the coordinated design of retroviral

vectors and packaging cell lines. The development of

packaging cell lines that package retroviral RNAs into

infectious particles without the concomitant production of

replication-competent virus created a new level of safety and

control. To do this, the structural genes necessary for

particle formation and replication, gag, pol, and env, were

integrated into cell lines without the RNA packaging signal,

psi+. Subsequent introduction of a retroviral vector

containing psi+, transcription and processing elements, and the gene of interest produces high-titer, replication-incompetent infectious virus. In other words, these retroviral particles can infect target cells and transmit the gene of interest, but cannot replicate within these cells since they lack the viral structural genes. The separate introduction and integration of the structural genes into the packaging cell line minimizes the chances of producing replication-competent virus due to recombination events during cell proliferation.

10

As a non-limiting, illustrative example, the following is presented. A Retro- X^{m} system is used to deliver and over-express antisense hNr-CAM gene in glioblastoma cells. Retro- X^{m} system is a complete retroviral gene expression system that can transduce up to 100% of cells.

- 15 Together with the RetroPackTM, PT67 cell line, the Retro-X Vectors produce infectious, replication-incompetent retrovirus that can be used to introduce a gene of interest into a wide variety of mammalian cell types in vitro or in vivoy. The highly efficient transduction machinery of
- 20 retroviruses can stably integrate the cloned gene into the host genome of nearly all mitotically dividing cells. A retroviral vector containing the gene of interest (hNr-CAM) is first transfected into the packaging cell line.
- Antibiotic selection can then be used to obtain a population of cells that stably expresses the integrated vector and, if desired, high-titer clones can be isolated from this population. Virus produced by either stably transfected cells can be used to infect target cells.

The hNr-CAM gene (first 1.3Kb) is cloned into the ECORI and BamHI site of the pLXSN retroviral vector in antisense direction. Human hNr-CAM specific primers (BT306,

^{5&#}x27; CATACGAATTCTAGATA CAACTAGTCTAATGCAGCTTAAAATAATGCC 3' SEQ.

ID. No.: 29; and BT307, 5' AGATAGATCCGCGGGATATCCATATT

CATTAGAGGCATTGGGATCCCATAC 3' SEQ. ID. No.: 30) are used to PCR amplify 1/3rd portion of the hNr-CAM gene. PCR is carried out using Gene Amp PCR kit from Perkin Elmer (Branchburg, NJ) under the following conditions: 4µl of dNTP mix, 2µl (100mg/µl) each of hNr-CAM specific primers, 4µl of 25µM MgCl₂, 125ng of cDNA template and 5 units of Amplitaq DNA polymerase. PCR amplification is carried out for 1 cycle at 94°C 3 min, 61°C 1 min, 72°C 4 min, then for 30 cycles at 94°C 1 min 61°C 1 min, 72°C 4 min followed by 1 cycle at 94°C 1 min, 61°C 1 min, 72°C 10 min.

10

The PCR product is run on a 1% agarose gel. band is then cut out and DNA was purified using gene clean system. DNA fragment is then digested with $5 \text{units}/\mu \text{l}$ EcoRI and BamHI for 2 hours at 37°C. Digested DNA is then run on a 2% agarose gel. The 1.3Kb fragment is then cut out, purified 15 and cloned in the antisense direction into EcoRI and BamHI sites of the pLXSN retroviral vector. Orientation of the cloned hNr-CAM gene is confirmed by sequencing. particular plasmids containing hNr-CAM anti-sense direction are termed "pLXSN-1/3Nr-AS". PLXSN and pLXSN-1/3Nr vectors 20 are transfected into Retropack™ PT67 cell line using the lipofectamine reagent. Cell lines are plated at a density of 3 x 104 cells/60mm. Twenty four hours after plating the cells are washed with serum-free media and transfected with lipofectamine reagent plus plasmid DNA (5 μ g) diluted in 1 ml of serum-free media. Cells are incubated at 37°C for 5 hours after which the reagent is replaced with media containing 10% FBS and cultures were incubated at 37°C for 72 hours. is then changed to an identical medium but containing adding 1000 $\mu g/ml$ G418 and incubated at 37°C for 96 hours. At this point, the medium is changed to one containing 200 μ g/ml Cultures are subsequently maintained in a medium containing 200 $\mu g/ml$ G418, changing the medium every 72-96 hours. Retroviral particles are then harvested by aspirating

the cell culture media into a sterile tube (approximately 10^5-10^6 recombinant virus particles/ml).

Viral titer is then determined using a protocol recommended by Clontech. Human glioblastoma cell lines (5GB, 1690-CRL, 1620-CRL) and two rat glioma cell lines C6 and 9L gliosarcoma cells are plated at a density of 3x10⁵ in two 100mm plates. Viral particles harvested from PT67 cell line culture are filtered through a 0.45µm filter and added on to human and rat glioma cell lines. Polybrene is added to a final concentration of 4µg/ml and incubated for 48 hours at 37°C. Cells are harvested and analyzed for hNr-CAM expression using Northern blot analysis. Cells that are expressing low levels of hNr-CAM are expanded in culture.

Northern blot analysis for the expression of hNr-CAM. Cell clones that are expressing low level of hNr-CAM are expanded in culture. Approximately 1x10⁷ glioblastoma cells 1690-CRL, 1620-CRL, HTB-16, C6, 9L gliosarcoma) will be injected subcutaneously into the flanks of ten female athymic nude mice (two sites each). Mice are then kept in a germ free environment. Tumor growth vill be analyzed every week for at least fourteen weeks and compared between anti-sense hNr-CAM and mock infected glioma cell lines.

It is appreciated that the physiological conditions under which a tumor develops in brain can be quite different from when it develops subcutaneously. As a non-limiting, illustrative example, the following is presented. The effect of hNr-CAM (antisense in 1690-CRL, 1620-CRL, HTB-16, C6, 9L cells) on tumor formation in brain can be assessed as described previously for a mutant EGFR (Nishikawa, et al., 1994, Proc. Natl. Acad. Sci. USA 91:7727-7731). Briefly, 5x10⁵ cells transfected or infected with the antisense hNr-30 CAM in 50-100µl of 1xPBS are inoculated into the cerebral hemisphere (using stereotectic instrument) of eight nude mice for each cell line. As a positive control for tumor growth,

U87MGΔEGFR cells that are known to form tumors in nude mice brains, can be used (Nishikawa, supra). 5x10⁵ U87MGΔEGFR cells are inoculated into the cerebral hemisphere of nude mice. Another group of nude nice are injected with pCMV-neo transfected or mock infected 1690-CRL, 1620-CRL, HTB-16, C6, 9L cells. Brains from all of these mice are removed at two week intervals, embedded in OCT compound, frozen in liquid nitrogen and stored at -80°C. 6μm sections are cut on a cryostat and immunocytochemistry are performed, for example, using the three markers described below.

10

GFAP (Glial Fibrillary Acidic Protein): Malignant glioma cells are known to cause gliosis of the surrounding tissue upon invasion (Mikkelsen, et al., 1998, Brain Tumor, Invasion, Bilogical, Clinical and Therapeutic Considerations, 15 Wiley Liss, N.Y.). Thus, the tissue area at the edge of originally transplanted glioblastoma cells and the mice brain tissue, can be stained by performing immunocytochemistry for the GFAP protein. After recovery of the brain tissue it is embedded in OCT blocks. Several $6\mu m$ sections are cut on a 20 cryostat. After washing sections in 1xPBS buffer, 200 μ l of diluted (1:80) rabbit anti-human GFAP are applied to slides. Slides are incubated for 18 hours at 4°C in a humid chamber. After washing in 1xPBS, FITC conjugated anti-rabbit immunoglobulins (1:50) (DAKO, A/S, Denmark) are applied and the slides are incubated at 24°C for 30 minutes in a humid 25 chamber. Cells are washed with 1xPBS and then stained with Hematoxylin (Richard Allen Scientific, Richland MI) for 30 seconds. Slides are then treated with a clarifying agent (Richard Allen Scientific, Richland MI) for 2 seconds and then in bluing agent (Richard Allen Scientific, Richland MI). 30 After washing in water, slides are coverslipped with 2% DABCO (Sigma, St. Louis MO) in 50% glycerol/lxPBS, and visualized with a Zeiss Axioskop UV microscope.

Cathepsin B (CB): CB is a cysteine protease that is expressed in gliomas directly in relation to grade of malignancy. CB is capable of degrading proteoglycans, a major component of the brain extracellular matrix, and could 5 be involved in the process of glial tumor cell invasion into peritumoral normal brain (Mikkelsen, supra). Each of the inoculated glioma cells (GB1690, C6, 9L cells transfected with pCMV-neo or pCMV-1/3Nr-AS) can examined for CB expression by performing immunocytochemistry for CB protein. It has been shown that the relative degree of granular stain 10 for glioma cell lines immunocytochemically in vivo correlates directly to the agree of invasiveness of the tumor displayed. To study the CB expression by immunocytochemistry, the same protocol as described above for the GEAP protein is followed. Human glioblastoma cell line U251MGn can be used as a 15 positive control for CB staining.

Ki67: Ki67 is a protein that is expressed at high levels in actively dividing cells (Mikkelsen, supra).
Immunocytochemistry can be performed on mouse brain sections

20 against the Ki67 antigen using a rabbit anti-human antibody.

The protocol for immunocytochemistry is the same as described above for the GFAP protein. U251MGn glioblastoma injected mice brain sections can be used as positive control for Ki67 specific staining. Direct comparison of the staining of these three markers provides the information about the extent of invasion of inoculated tumor cells.

8. EXAMPLE: IDENTIFICATION OF GENES ALTERED BY hnr-cam

In order to identify genes that are altered by the hNr-CAM gene product in 5GB gliioblastoma cells, we compared the expression of 5000 genes in pCMV-neo or pCMV-1/3Nr-AS transfected 5GB glioblastoma cells using the Array technique. Two identical Human GeneFilters™ were differentially

hybridized with cDNA prepared from pCMV-neo or pCMV-1/3Nr-AS transfected 5GB glioblastoma cells. Two identical array membranes containing 5000 genes were purchase dfrom REsearch Genetics. The membranes were prehyridized in a pre5 hybridization solution for 12 hours. Hybridization was done with a 1x105 cpm/ml cDNA probe. The probe was prepared by carrying out 1st strand synthesis from pCMV-neo or pCMV-1/3Nr-AS transfected 5GB glioblastoma cells 1µg polyA+mRNA. First strand cDNA synthesis was carried out using the Advantage cDNA synthesis kit from Clontech. The membranes were washed in a wash solution (0.1%SDS/1XSSC) for 30 minutes at room temperature and then at 50°. Membranes were then exposed to X-ray film. Results are presented in Figures 26 (A and B).

As shown in Figures 26 (A and B), two genes were identified that were differentially expressed. Selectin (endothelial adhesion molecule 2) was detected in pCMV-neo transfected cells (Figure 26B) and not n pCMV-1/3Nr-AS (Figure 26A) transfected 5GB cells. A novel gene (accession # H7785) was detected in pCMV-1/3Nr-AS (Figure 26A) transfected in pCMV-neo transfected (Figure 26B) 5GB cells. We are not only interested in exploring the role of these genes in glioblastoma cells in the context of hNr-CAM over-expression but also inunderstanding the mechanism by which hNr-CAM modulates the expression of these genes.

25
The present invention is not to be limited in scope by the microorganism deposited or the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

PCT/US99/09039

WO 99/55380

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

5

10

15

20

25

30

WHAT IS CLAIMED IS:

 A pharmaceutical composition for the inhibition or treatment of tumorigenesis comprising an antisense nucleic acid complementary to at least a portion of an RNA transcript of a Nr-CAM gene in an amount effective to inhibit hyperproliferation of a tumor cell.

- 2. A pharmaceutical composition for the inhibition or treatment of tumorigenesis comprising an antibody to Nr-CAM in an amount effective to inhibit hyperproliferation of a tumor cell.
- 3. A method of treating, inhibiting or preventing a disease or disorder involving cell overproliferation in a subject comprising administering to a subject in which such treatment or prevention is desired an effective amount of a molecule that inhibits Nr-CAM function.
- 4. The method according to claim 3 in which the disease or disorder is a malignancy.
 - 5. The method according to claim 3 in which the disease or disorder is selected from the group consisting of brain cancer, leukemia and B cell lymphoma.
 - 6. The method according to claim 3 in which the subject is a human.
- 7. The method according to claim 5 in which the 30 brain cancer is selected from the group consisting of glioblastoma, glioma, meningioma, astrocytoma, medulloblastoma, neuroectodermal cancer and neuroblastoma.

8. The method according to claim 7 in which the glioblastoma is glioblastoma multiforme.

- 9. The method according to claim 3 in which the disease or disorder is selected from the group consisting of premalignant conditions, benign tumors, hyperproliferative disorders, and benign dysproliferative disorders.
- molecule that inhibits Nr-CAM function is selected from the group consisting of an anti-Nr-CAM antibody or a fragment thereof, a Nr-CAM derivative or analog that is capable of being bound by an anti-Nr-CAM antibody, a Nr-CAM antisense nucleic acid, and a nucleic acid comprising at least a portion of a Nr-CAM gene into which a heterologous nucleotide sequence has been inserted such that said heterologous sequence inactivates the biological activity of at least a portion of the Nr-CAM gene, in which the Nr-CAM gene portion flanks the heterologous sequence so as to promote homologous recombination with a genomic Nr-CAM gene.

20

25

- 11. The method according to claim 3 in which the molecule that inhibits Nr-CAM function is an oligonucleotide which (a) consists of at least six nucleotides; (b) comprises a sequence complementary to at least a portion of an RNA transcript of a Nr-CAM gene; and (c) is hybridizable to the RNA transcript under moderately stringent conditions.
- 12. A method of inhibiting, treating or preventing a disease or disorder involving cell proliferation in a subject comprising administering to a subject in need of such treatment an effective amount of a molecule that promotes Nr-CAM function.

13. The method according to claim 12, in which the disease or disorder is selected from the group consisting of degenerative disorders, growth deficiencies, hypoproliferative disorders, physical trauma, lesions, and wounds.

14. A method of diagnosing a disease or disorder characterized by an aberrant level of Nr-CAM RNA or protein in a subject, comprising measuring the level of Nr-CAM RNA or protein in a sample derived from the subject, in which an increase or decrease in the level of Nr-CAM RNA or protein, relative to the level of Nr-CAM RNA or protein found in an analogous sample from another subject not having the disease or disorder, indicates the presence of the disease or disorder in the subject.

15

- 15. A method of diagnosing or screening for the presence of or a predisposition for developing a disease or disorder involving cell overproliferation in a subject comprising detecting Nr-CAM DNA, RNA or protein derived from the subject in which the presence of said Nr-CAM DNA, RNA or protein indicates the presence of the disease or disorder or a predisposition for developing the disease or disorder.
- molecule selected from the group consisting of an anti-Nr-CAM antibody, a nucleic acid probe capable of hybridizing to a Nr-CAM RNA, a pair of nucleic acid primers capable of priming amplification of at least a portion of a Nr-CAM nucleic acid, an anti-Nr-CAM ligand antibody, a nucleic acid probe capable of hybridizing to an Nr-CAM ligand encoding RNA and a pair of nucleic acid primers capable of priming amplification of at least a portion of an Nr-CAM ligand encoding nucleic acid.

17. A pharmaceutical composition for the inhibition or treatment of tumorigenesis comprising an antisense nucleic acid complementary to at least a portion of an RNA transcript of an Nr-CAM ligand encoding gene in an amount effective to inhibit hyperproliferation of a tumor cell.

- 18. A pharmaceutical composition for the inhibition or treatment of tumorigenesis comprising an antibody to an Nr-CAM ligand encoding gene in an amount effective to inhibit hyperproliferation of a tumor cell.
- 19. A method of inhibiting, treating or preventing a disease or disorder involving cell overproliferation in a subject comprising administering to a subject in which such treatment or prevention is desired an effective amount of a molecule that inhibits an Nr-CAM ligand encoding gene function.
- characterized by an aberrant level of an Nr-CAM ligand encoding RNA or protein in a subject, comprising measuring the level of an Nr-CAM ligand encoding RNA or protein in a sample derived from the subject, in which an increase or decrease in the level of said RNA or protein, relative to the level of said RNA or protein found in an analogous sample from another subject not having the disease or disorder, indicates the presence of the disease or disorder in the subject.
- 21. A method of diagnosing or screening for the 30 presence of or a predisposition for developing a disease or disorder involving cell overproliferation in a subject comprising detecting an Nr-CAM ligand encoding DNA, RNA or

protein derived from the subject in which the presence of said DNA, RNA or protein indicates the presence of the disease or disorder or a predisposition for developing the disease or disorder.




FIGURE 1

181 ctgtgccaga tgattagtgc actggaagta cctcttgatc caasacttct tgaagaottg 241 gtacagcete caaccatcae ccaacagtet ccaaaagatt acattattga ccctegggag 301 aatattgtaa tooagtgtga agccaaaggg aaaccgcccc caagetttte ctggacccgt 361 aatgggaete attttgacat egataaagae eetetggtea eeatgaagee tggcaeagga

421 acgctcataa ttaacatcat gagcgaaggg aaagctgaga cctatgaagg agtctatcag
481 tgtacagcaa ggaacgaacg cggagctgca gtttctaata acattgttgt ccgcccatcc
541 agatcaccat tgtggaccaa agaaaaactt gaaccaatca cacttcaaag tggtcagtot
601 ttagtacttc cctgcagacc cccaattgga ttaccaccac ctataatatt ttggatggat 661 aatteette aaagaettee acaaagtgag agagtteete aaggtttgaa tggggaeett 721 tatttteea atgteeteee agaggaeaee egegaagaet atatetgtta tgetagattt

781 aatcatactc aaaccataca gcagaagcaa cctattetg tgaaggtgat ttcagtggat 841 gaattgaatg acactatagc tgotaatttg agtgacactg agttttatgg tgctaaatca 901 agtagagaga ggccaccaac atttttaact ccagaaggca atgcaagtaa caaagaggaa 961 ttaagaggaa atgtgettte actggagtge attgeagaag gactgeetae eccaattatt
1021 taetgggeaa aggaagatgg aatgetaeee aaaaacagga cagtttataa gaactttgag 1081 aaaaccttgc agatcattca tgtttcagaa gcagactctg gaaattacca atgtatagca 1141 aasaatgcat taggagccat ccaccatacc atttetetta gagttaaagc ggctccatac 1201 tggatcacag cccctcaaaa tettgtgetg tecccaggag aggatgggac ettgatetge 1261 agagetaatg geaaceccaa acccagaatt agetggttaa caaatggagt eccaatagaa

1321 attgeectg atgaceccag cagaaaaata gatggegata caaarggagt cecaarayaa
1321 attgeectg atgaceccag cagaaaaata gatggegata caatgtttt tteaaatgtt
1331 caagaaagat caagtgeagt atateagtgelaatgeeteta atgatatatg atalltaetg
1441 geaaacgeat ttgtaaatgt getggetgag ceaecaegaa teeteaace tgaaaacaec
1501 etetaecagg teattgeaaa caggeetget ttaetagaet gtgeettett tgggteteet

1561 eteccaacca tegagtggtt taaaggaget aaaggaagtg etetteatga agatatttat 1621 gttttacatg aaaatggaac tttggaaatc aaagatgota catggatcgt taaagaaatt 1681 cctgtggccc aaaaggacag tacaggaact tatacgtgtg ttgcaaggaa taaattaggg 1741 atggcaaaga atgaagttca cttacagocc gaatatgcag ttgtgcaaag agggagcatg
1801 gtgtcctttg aatgcaaagt gaaacatgat cacaccttat ccctcactgt cctgtggctg
1861 aaggacaaca gggaactgcc cagtgatgaa aggttcactg ttgacaagga tcatctagtg 1921 gtagetgatg teagtgaega tgacageggg acetacaegt gtgtggeeaa caccactetg

1981 gacagegtet cegecagege tgtgettage gttgttgete etactecaae tecageteee 2041 gtttacgatg teccaaatec tecetttgae ttagaactga cagateaact tgacaaaagt 2101 gttcagetgt catggacece aggegatgae aacaatagee ceattacaaa attcateate 2161 gaatatgaag atgcaatgca caagccaggg ctgtggcacc accaaactga agtttctgga 2221 acacagaeca cageceaget gaagetgtet cettaegtga actaeteett eegegtgatg 2281 gcagtgaaca gcattgggaa gagcttgccc agogaggcgt ctgagcagta tttgacgaaa 2341 gcctcagaac cagataaaaa ccccacagct gtggaaggac tgggatcaga gcctgataat 2401 ttggagatta cgtggaagcc cttgaatggt ttcgaatcta atgggccagg ccttcagtac

2461 aaagttaget ggegecagaa agatggtgat gatgaatgga catetgtggt tgtggcaaat 2521 gtatecaaat atattgtete aggeacgoca acetttgtte catacetgat caaagtteag 2581 geectgaatg acatggggtt tgcececgag ceagetgtag teatgggaca ttetggagaa 2641 gacctcccaa tggtggctcc tgggaaogtg cgtgtgaatg tggtgaacag tacattagcc 2701 gaggtgcact gggacccagt acctctgaaa agcatccgag gacacctaca aggctatogg 2761 atttactatt ggaagaccca gagttcatct aasagaaaca gacgtcacat tgagaaaaag 2821 atceteacet tecaaggeag caagacteat ggeatgttge eggggetaga gecetttage 2881 cactacacac tgaatgteeg agtggteaat gggaaagggg agggeeage cagecetgae 2941 agagtetta atactecaga aggagteece agtgeteect egtetttgaa gattgtgaat 3001 ccaacactg acteteteac titggaatgg gatecacoga gecaccogaa tggcatittg 3061 acagagtaca cettaaagta teagecaatt aacagcacao atgaattagg ccetetggta

3121 gatttgaaaa ttcctgccaa caagacacgg tggactttaa aaaatttaaa tttcagcact 3181 cyatataagt titatitota tyoacaaaca toagoaggat oaggaagtoa aattacagag 3241 gaagoagtaa caactytyga tyaagotygt attottooac cigatytagy tyoaggoaaa 3241 gaageageaa caactycyya cyaayctygt attottooae ctgatgtagy tycayycaaa 3301 gttcaagetg taaataccag gatcagcaat cttactgctg cagetgctga gacctatgoc 3361 aatatcagtt gggaatatga gggaccagag catgtgaact tttatgttga atatggtgta 3421 gcaggcagca aagaagaaty gagaaaagaa attgtaaaty gttotoggag cttctttggg 3481 ttaaagggtc taatgccagg aacagcatac aaagttogag ttggtgctgt gggggactot 3541 ggttttgtga gttcagagga tgtgtttgag acaggocoag ogatggcaag coggoaggtg 3601 gatattgcaa ctcagggctg gttcattggt ctgatgtgt ctgttgctct ccttatotta

3661 attitgetga tigtitgett cateagaga accaegaga accaegagitg giaaatatee agitaaagaa 3721 aaggaagatg cecateetga eeetgataate cageotatga aggaagatga tgggacattt 3841 gacaggactg tgaaaaaaga agacacacaag cettgaaaa aaggaagteg aacteettea 3841 gacaggactg tgaaaaaaga agatagtga gacagcetag ttgacaatg agaaggggtt 3901 aatggecagt teaatgaga tggeteettt attggacaat acagtggtaa gaaagagaaa 3861 gacagagata aagagaaaa aaggagatga agaagagaaa aagagagaa acaagagaaa aagaggaatga acaagagaaa aagagagaa aagagagaa aagaggaatgaa aagagagaa

3961 gagcoggetg aaggaaacga aagctcagag gcacettote etgteaacge catgaattee 4021 tttgtttaat ttttaagete aaagccaata ttggatttet etagaatgt tateetaage 4081 tettgtttgt cagceetete atactatgaa valgtgggta gagagtatat ttte

FIGURE 2A


BT180 BTITE


"MPKKGRLSAGRVPLILFLCOMISALEVPLDPKLLEDLVQPPTIT
QQSPKDYIIDPRENIVIQCEAKGKPPPSFSWTRNGTHFDIDKDPLVTMKPGTGTLIIN
IMSEGKAETYEGVYQCTARNERGAAVSNNIVVRPSRSPLWTKEKLEPITLQSGQSLVL
PCRPPIGLPPPIIFWMDNSFQRLPQSERVSQGINGDLYFSNVLPEDTREDYICYARFN
HTQTIQQKQPISVKVISVDELNDTIAANLSDTEFYGAKSSRERPPTFLTPEGNASMKE
ELRGNVLSLECLAEGLPTPIIYWAKEDGMLPKNRTVYKNFEKTLQIIHVSEADSGNYQ
CIAKNALGAIHHTISVRVKAAPYWITAPQNLVLSPGEDGTLICRANGNPKPRISWIMN
GVPIELAPDDPSRKIDGDTIIFSNVQERSSAVYQCNASNEYGYLLANAFVNVLAEPPR
ILTPANTLYQVIANRPALLDCAFFGSPLPTIEWFKGAKGSALHEDIYVLHENGTLEIK
DATWIVKEIPVAQKDSTGTYTCVARNKLGMKNEVHLQPEYAVVQRGSMVSFECKVKH
DHTLSLTVIMLKDNRELPSDERFTVDKDHLVVADVSDDDSGTYTCVANTTLDSVSASA
VLSVVAPTPTPAPVYDVPNPPFDLELTDQLDKSVQLSWTPGDDNNSPITKFIIEYEDA
MRKPGLWHEQTEVSGTQTTAQLKLSPYVNYSFRVMAVNSIGKSLESEASEQYLTKASE
PDKNPTAVEGLGSEPDNLEITWKPLNGFESNGPGLQYKVSWRQKDGDDEWTSVVVANV
SKYIVSGTPTFVPYLIKVQALNDMGFAPEPAVVMGHSGGDLPMVAPGNVRVNVNSTL


AEVHWDPVPLKSIRGHLQGYRIYYWKTQSSSKRNRRHIEKKILTFQGSKTHGALPGIE PFSHYTLNVRVVNGKGEGPAS PDRVFNTPEGVPSAPSSLKIVNPTLDSL/TLEMDPPSH PNGILTEYTLKYQPINSTHELGPLVDLKI PANKTRWTLKNINFSTRYKFYFYAQTSAG SGSQITEEAVTTVDEAGILPPDVGAGKVQAVNTRISNLTAAAAETYANISWEYEGPEH VNFYVEYGVAGSKEEWRKEIVNGSRSFFGLKGLMPGTAYKVRVGAVGDSGFVSSEDVF ETGPAMASRQVDIATQGWFIGIACAVALLILILLLIVCFIRRNKGGKYPVKEKEDAHAD PEIQPMKEDDGTFGEYSDAEDHKPLKKGSRTPSDRTVKKEDSDDSLVDYGEGVNGQFN EDGSFIGQYSGKKEKEPAEGNESSEAPSPVNAMNSFV**


HUMAN NT-CAM	4097	TCTCATACTATGAACATATGGGTAGAGAGTATATTTTC *****************************	4134
D4-1	1	TCTCATACTATGAACATATGGGTAGAGAGTATATTTTCTGCTGT	44
RAT Nr-CAM	3713	TCTCATACTATGGACATATGGGTAGAAAGAATGTTTTCTGOGGT	3 757
D4-1	45	ATGTTAGTATTATGAGAATAGTTACAGCAAAAACATAACTCAGT	89
RAT NI-CAM	3758	ATATGAGTATTATAAGAACAGAGCAAGAACATAACTCAGTCAG	3802
D4-1	90	CRAAGTATATGTTAATATGAACTGGAATGCAAAAGTGCATACTT	134
RAT Nr-CAM	3802	CAGATGATACGTTAATATGAACTGGGGTGAAAAGG	383:
D4-1	135	AAAAAAAAAAA TAATII AAAAAAAAAAAAAAAAAAAA	179

hNr-CAM hNr-CAM1/3AS













Cytoplasmic tail


FIGURE 2D




FIGURES 3A-F




FIGURES 4A-B




FIGURES 5A-B



FIGURES 6A-B



FIGURES 7A-B



FIGURES 8A-B





FIGURE 9






FIGURE 11




FIGURE 12





FIGURE 13






FIGURE 14A-B




FIGURE 15A-D




FIGURE 16A-D




FIGURE 17




FIGURE 18





FIGURE 19






FIGURE 20A-C




FIGURE 21




FIGURE 22A





FIGURE 22B






FIGURE 23A-C

	. and securit at att the	
Ruman Hr-CAK: Rat Hr-CAK:	1 ATGCCCALADAMACCCCTATCTCCCCCCCCAAAAAAAAAAAAAAAAAA	•
Ruman Mr-CM:	61 ATOATTAGTOCACTOGRAGIAGICATGTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT	
RAT MI-CAN:	79 ATGATOAGGETCTGGATGTTCTTGATGTACTTGATGTACTTGGTACAGGCT 138	
	121 CCACCATCACCCACAGTCTCCALAMGATTACATTATTGACCCTCGGGGGGAATATTGTA 160	
	DIS CONTRACTORISMONTOLOGICALIS	
	181 ATCCMOTOTOLAGCCAAAGGGAAACGCCCCCAAGCTTTTCCTGGACCCOTAATGGGACT 240	
	THE WICH ALL PROPERTY OF THE PROPERTY SEED AND THE PROPERTY SEED A	,
	241 CATTITUDATEGRADAGACCTETOGICACCATORAGCTGGGAACCCTGATA 300	
	301 ATTACATCATGAGGGAAGGGGAAAGCTGAGAGGTTATGAAGGAGGTTATGAGGGGAGGGA	
	361 ACCIACOMACCICOMOTICICACIATACATICITATION (100)	
Rat Mr-CAN:	1111 11 111111111 11 11 11111111111111	
	421 TTOTGGACCHAGANAACTTGAACCACTTCAAGTGGTCATCTTTAGACTT 480	
	175 THE CONTRACT OF THE PROPERTY AND ADDRESS OF THE PROPERTY ADDRESS OF THE PR	
Rat Er-CAN:	461 CCCTGCAGACCCCCAATTGGATTACCACCACCATAATATTTTTGGATGGA	
Human Hr-CAK:	499 CCATGTAGGCCTCCAATTGGATTACCACCGGCCATAATATTTTGGATGGA	
RAT Mr-CM:	SSP CANAGACTOCCACAGAGTAGAGCOGGTTCCCCAAGGACTGAATGGAGACCTTTACTTCTCC 618	
Ruman Fr-CAR:	COL MATORICITICO MAGRACIA CONCOMMINA CITATA DE TRANSPORTA COMPANSA DE CASA	
Rat Mr-CAM:	11111111111111111111111111111111111111	
Busan Nr.CAM:	661 CAMCCATACAGCAGAAGCAACCAATTCAGTGAAAGGGGAATGAAT	
Rat Mr-CAM:	738	
RAT Mr-CM:	721 GACACTATAGCTGCTAATTTUAGTGCACTGAGTTTTATGGTGCTAATCAAGTAGAAGG 780	
Ruman Fr-CAK:	781 AGGCCACCACATTITTAACTCCAGAAGGCAATGCAAGTTAACTAAGTAAGTAAG	
Rat Br-CNN:	799 AGGCCACCALCATTICTALCTCCAGAGGCCATGALAGTCACALGGAAGTALTALGAGGA 858	
Human Hr-CAN:	841 AATGIGCTITCACTGGAGTGCATTGCAGAAGTACTTCCAGAAAGTACTTCCAGAAGTACTTCCAGAAGTACTTCCAGAAGTACTCAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGTACTACAAGAAGAAGTACTACAAGAAGAAGTACTACAAGAAGAAGAAAGA	
Rat Mr-CMI	11	
Human Nr-CAN:	901 MOGNAGATOGINTOCTICOCCHINICAGACAGTTATAGACACTTICAGAMACATTG 960	
Rat Hr-CM:	919 AAGGAMATGGBACOCTTCCTUTCAACCGBACGTTTTATCGGAACCTTGAGAAACCTTG 978	
hman Hr-CAR:	961 CAGATCATTCATGUTTCAGAAGCAGACTCTGGAAATTACCAATGTATACCAAAAAATGCA 1020	
Rat Hr-CAN:	979 CHARCATTCATGTCTCTGAAGCAGACTCTGGAAATTATCAGTGCATAGCAAAAAAAGCA 1038 1021 TAGGAGCCATCCACCATACCATTCTGTTAGAGTTAAAGGGGCCCCATACTGGATCACA 1088	
Rat Mr-CAN:	1039 TTOOGREGOTCCATCCATACCATTCTCACCATTAAAGCCCCCCCACTGGATTGTT 1098	
man M-CH.		
RAT Nr-CAH:	1099 GOCCTONONCCTOTICCCCONGGUNDARGGUNCCCCCATCTOCHARGCTANC 1158	
Burn Hr-CM:	1141 GGCALCCCALACTAGCTGGTTACAAATGGAGTGCCAATAGAAATGCCCCT 1200	FIGL
Rat Hr-CAN:	1159 GCCACCCAAACCCAGAATAGCTGGTTAACAAATGGAGTGCCAGTAGAAATGCTCTC 1218	
Human Hr-CAK:	1201 GANGACCCAGAAAAAAAAGGCGATACCATTATATTATAAANOTAAAAAAA 1260 	24
Banan Mr-CM	1261 TOMOTOCOTATATOCOTOCATOCATATAMATATATATATATOCOMACOCA 1120	
RAE RY-CHI	1279 TOMOROWITATOMOCATOCATORATEROWATETACTMOCATEGAL 1318	
Busan Hr-CAK	: 1121 TITUTANATOTOCTOMOCACCACANATOCTCACACCTOCANCACA 1271	
Bat Er-Car.	1110 THE THE PROPERTY OF THE PR	

URE




FIGURE 25A

FIGURE 25B

pCMV-1/3Nr-AS



FIGURE 26A

pCMV-neo



FIGURE 26B

SEQUENCE LISTING

.<110> PACIFIO	NORTHWEST	רבאורבים יי	רוואווי א מינאווי	NT.			
, CIIV FACILI	C NORTHWEST	CANCER F	CONDATIO	N			
<120> Nr-CAM FOR THI	GENE, NUCLE ERAPEUTIC AN	IC ACIDS D DIAGNO	AND NUC	LEIC ACI S FOR TU	D PRODUC MORS	TS	
<130> 8511-0	21-228		•				
<140> PCT/US	99/						
<141> 1999-0							
<150> 60/112							
<151> 1998-1	2-14		-				
<150> 60/083	160						
<151> 1998-0				•			
, , , , , , , , ,				-			
<160> 32							
<170> Patent	In Ver. 2.0						
<210> 1							
<211> 4134							
<212> DNA							
<213> Homo s	apiens						
<220>							
<221> CDS						•	
<222> (130).	. (4029)						
	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,						
<400> 1							
cttcaaagtt c	cccgcatga aa	attactta	aacgtto	cac acaa	egtttc a	gaaaatctt	60
ttgtgaaaga a	gaaaaggaa at	tcagtgtg	g tgagtct	cag cag	gagttaa g	gctaatgcag	120
cttaaaata at	g ccg aaa aa	ag aag co	gc tta to	t gcg gg	gc aga gt	g ccc ctg	171
Me	t Pro Lys Ly	/s Lys Aı	rg Leu Se	er Ala G	ly Arg Va	al Pro Leu	
	1	5		;	10	-	
att ctc ttc	cta tac cea	atg att	agt ggs	cta ass	ata	at =	210
Ile Leu Phe	Leu Cys Gln	Met Ile	Ser Ala	Leu Glu	Val Pro	Leu Aen	219
15	20			25	var Pro	30	
cca aaa ctt	ctt gaa gac	ttg gta	cag cct	cca acc	atc acc	caa cag	267
Pro Lys Leu		Leu Val	Gln Pro	Pro Thr	Ile Thr	Gln Gln	
	35		40			45	
tct cca aaa	gat tac att	att dec	cct cc-	G=G	3++		775
Ser Pro Lys	Asp Tvr Ile	Ile Asp	Pro Arg	Glu Asp	Tle Val	atc cag	315
	50	P	55	and Wall	60	TIE GIN	
tgt gaa gcc	aaa ggg aaa	ccg ccc	cca agc	ttt tcc	tgg acc	cgt aat	363
Cys Glu Ala	rive Gly rive		Pro Ser	Phe Ser		Arg Asn	
65		70			75		

																•
Gly 999	act Thr 80	cat His	ttt Phe	gac Asp	atc Ile	gat Asp 85	aaa Lys	gac Asp	cct Pro	ctg Leu	gtc Val 90	acc Thr	atg Met	aag Lys	cct Pro	411
ggc Gly 95	aca Thr	gga Gly	acg Thr	ctc Leu	ata Ile 100	att Ile	aac Asn	atc Ile	atg Met	agc Ser 105	gaa Glu	ggg Gly	aaa Lys	gct Ala	gag Glu 110	459
acc Thr	tat Tyr	gaa Glu	gga Gly	gtc Val 115	tat Tyr	cag Gln	tgt Cys	aca Thr	gca Ala 120	agg Arg	aac Asn	gaa Glu	cgc Arg	gga Gly 125	gct Ala	507
gca Ala	gtt Val	tct Ser	aat Asn 130	aac Asn	att Ile	gtt Val	gtc Val	cgc Arg 135	cca Pro	tcc Ser	aga Arg	tca Ser	cca Pro 140	ttg Leu	tgg Trp	555
acc Thr	aaa Lys	gaa Glu 145	aaa Lys	ctt Leu	gaa Glu	cca Pro	atc Ile 150	aca Thr	ctt Leu	caa Gln	agt Ser	ggt Gly 155	cag Gln	tct Ser	tta Leu	603
gta Val	ctt Leu 160	ccc Pro	cys	aga Arg	ccc Pro	cca Pro 165	att Ile	gga Gly	tta Leu	cca Pro	cca Pro 170	cct Pro	ata Ile	ata Ile	ttt Phe	651
tgg Trp 175	atg Met	gat Asp	aat Asn	tcc Ser	ttt Phe 180	caa Gln	aga Arg	ctt Leu	cca Pro	caa Gln 185	agt Ser	gag Glu	aga Arg	gtt Val	tct Ser 190	699
caa Gln	ggt Gly	ttg Leu	aat Asn	999 Gly 195	gac Asp	ctt Leu	tat Tyr	ttt Phe	tcc Ser 200	aat Asn	gtc Val	ctc Leu	cca Pro	gag Glu 205	gac Asp	747
acc Thr	cgc Arg	gaa Glu	gac Asp 210	tat Tyr	atc Ile	tgt Cys	tat Tyr	gct Ala 215	aga Arg	ttt Phe	aat Asn	cat His	act Thr 220	caa Gln	acc Thr	795
ata Ile	cag Gln	cag Gln 225	aag Lys	caa Gln	cct Pro	att Ile	tct Ser 230	gtg Val	aag Lys	gtg Val	att Ile	tca Ser 235	gtg Val	gat Asp	gaa Glu	843
ttg Leu	aat Asn 240	gac Asp	act Thr	ata Ile	gct Ala	gct Ala 245	aat Asn	ttg Leu	agt Ser	gac Asp	act Thr 250	gag Glu	ttt Phe	tat Tyr	ggt Gly	891
gct Ala 255	aaa Lys	tca Ser	agt Ser	aga Arg	gag Glu 260	agg Arg	cca Pro	cca Pro	aca Thr	Phe 265	Leu	act Thr	cca Pro	gaa Glu	ggc Gly 270	9 39
aat Asn	gca Ala	agt Ser	aac Asn	aaa Lys 275	gag Glu	gaa Glu	tta Leu	aga Arg	gga Gly 280	Asn	gtg Val	ctt Leu	tca Ser	Ctg Leu 285	Glu	987
tgc Cys	att	gca Ala	gaa Glu _290	Gly	ctg Leu	cct Pro	acc Thr	cca Pro _295	Ile	att	tac Tyr	tgg Trp	gca Ala _300	Lys	gaa Glu	1035

						•											
g A	at sp	gga Gly	atg Met 305	cca Leu	ccc Pro	aaa Lys	aac Asn	agg Arg 310	aca Thr	gtt Val	tat Tyr	aag Lys	aac Asn 315	ttt Phe	gag Glu	aaa Lys	1083
T	cc hr	ttg Leu 320	cag Gln	atc Ile	att Ile	cat His	gtt Val 325	tca Ser	gaa Glu	gca Ala	gac Asp	tct Ser 330	Gly	aat Asn	tac Tyr	caa Gln	1131
C	gt ys 35	ata Ile	gca Ala	aaa Lys	aat Asn	gca Ala 340	tta Leu	gga Gly	gcc Ala	atc Ile	cac His 345	cat His	acc Thr	att Ile	tct Ser	gtt Val 350	1179
a A	ga Irg	gtt Val	aaa Lys	gcg Ala	gct Ala 355	cca Pro	tac Tyr	tgg Trp	atc Ile	aca Thr 360	gcc Ala	cct Pro	caa Gln	aat Asn	ctt Leu 365	gtg Val	1227
I	tg .eu	tcc Ser	cca Pro	gga Gly 370	gag Glu	gat Asp	61 y	acc Thr	ttg Leu 375	atc Ile	tgc Cys	aga Arg	gct Ala	aat Asn 380	ggc Gly	aac Asn	1275
F	ro	a aa Lys	ccc Pro 385	aga Arg	att Ile	agc Ser	tgg Trp	tta Leu 390	aca Thr	aat Asn	gga Gly	gtc Val	cca Pro 395	ata Ile	gaa Glu	att Ile	1323
Į	gcc Ala	cct Pro 400	gat Asp	gac Asp	ccc Pro	agc Ser	aga Arg 405	aaa Lys	ata Ile	gat Asp	ggc Gly	gat Asp 410	acc Thr	att Ile	att Ile	ttt Phe	1371
5	ca Ser	aat Asn	gtt Val	caa Gln	gaa Glu	aga Arg 420	tca Ser	agt Ser	gca Ala	gta Val	tat Tyr 425	cag	tgc Cys	aat Asn	gcc Ala	tct Ser 430	1419
i	aat Asn	gaa Glu	tat Tyr	gga Gly	tat Tyr 435	Leu	ctg Leu	gca Ala	aac Asn	gca Ala 440	ttt Phe	gta Val	aat Asn	gtg Val	ctg Leu 445	gct Ala	1467
. (gag Glu	cca Pro	cca Pro	cga Arg 450	Ile	ctc Leu	aca Thr	cct Pro	gca Ala 455	aac Asn	aca Thr	ctc Leu	tac Tyr	cag Gln 460	gtc Val	att Ile	1515
1	gca Ala	aac Asn	agg Arg 465	Pro	gct Ala	tta Leu	Leu	gac Asp 470	Cys	gcc Ala	ttc Phe	ttt Phe	999 Gly 475	tct Ser	cct Pro	ctc Leu	1563
1	cca Pro	acc Thr 480	Ile	gag Glu	tgg Trp	ttt Phe	aaa Lys 485	gga Gly	gct Ala	aaa Lys	gga Gly	agt Ser 490	gct Ala	ctt Leu	cat His	gaa Glu	1611
4	gat Asp 495	att	tat Tyr	gtt Val	tta Leu	cat His 500	Glu	aat Asn	gga Gly	act Thr	ttg Leu 505	gaa Glu	atc Ile	aaa Lys	gat Asp	gct Ala 510	1659
	aca Thr	tgg Trp	ato	gtt Val	aaa Lys 515	Glu	att Ile	cct Pro	gtg Val	gcc Ala 520	Gln	aag Lys	gac Asp	agt Ser	aca Thr 525	gga Gly	1707

WO 99/55380 PCT/US99/09039 .

																•
act Thr	tat Tyr	Thr	tgt Cys 530	gtt, Val	gca Ala	agg Arg	aat Asn	aaa Lys 535	tta Leu	ggg Gly	atg Met	gca Ala	aag Lys 540	aat Asn	gaa Glu	1755
gtt Val	cac His	tta Leu 545	cag Gln	ccc Pro	gaa Glu	tat Tyr	gca Ala 550	gtt Val	gtg Val	caa Gln	aga Arg	999 Gly 555	agc Ser	atg Met	gtg Val	1803
tcc Ser	ttt Phe 560	gaa Glu	tgc Cys	aaa Lys	gtg Val	aaa Lys 565	cat His	gat Asp	cac His	acc Thr	tta Leu 570	tcc Ser	ctc Leu	act Thr	gtc Val	1851
ctg Leu 575	tgg Trp	ctg Leu	aag Lys	gac Asp	aac Asn 580	agg Arg	gaa Glu	ctg Leu	ccc Pro	agt Ser 5 85	gat Asp	gaa Glu	agg Arg	ttc Phe	act Thr 590	1899
gtt Val	gac Asp	aag Lys	gat Asp	cat His 595	cta Leu	gtg Val	gta Val	gct Ala	gat Asp 600	gtc Val	agt Ser	gac Asp	gat Asp	gac Asp 605	agc Ser	1947
99 9 Gly	acc Thr	tac Tyr	acg Thr 610	tgt Cys	gtg Val	gcc Ala	aac Asn	acc Thr 615	act Thr	ctg Leu	gac Asp	agc Ser	gtc Val 620	tcc Ser	gcc Ala	1995
agc Ser	gct Ala	gtg Val 625	ctt Leu	agc Ser	gtt Val	gtt Val	gct Ala 630	cct Pro	act Thr	cca Pro	act Thr	cca Pro 635	gct Ala	ccc Pro	gtt Val	2043
tac Tyr	gat Asp 640	gtc Val	cca Pro	aat Asn	cct Pro	ccc Pro 645	ttt Phe	gac Asp	tta Leu	gaa Glu	ctg Leu 650	aca Thr	gat Asp	caa Gln	ctt Leu	2091
gac Asp 655	aaa Lys	agt Ser	gtt Val	cag Gln	ctg Leu 660	tca Ser	tgg Trp	acc Thr	cca Pro	ggc Gly 665	gat Asp	gac Asp	aac Asn	aat Asn	agc Ser 670	2139
ccc Pro	att Ile	aca Thr	aaa Lys	ttc Phe 675	Ile	atc Ile	gaa Glu	tat Tyr	gaa Glu 680	gat Asp	gca Ala	atg Met	cac His	aag Lys 685	Pro	2187
ggg Gly	ctg Leu	tgg	cac His 6 90	His	caa Gln	act Thr	gaa Glu	gtt Val 695	Ser	gga Gly	aca Thr	cag Gln	acc Thr 700	Thr	gcc Ala	2235
caç Glr	ctg Leu	aag Lys 705	Leu	tct Ser	cct Pro	tac Tyr	gtg Val 710	Asn	tac Tyr	tcc Ser	ttc Phe	cgc Arg 715	Val	atg Met	gca Ala	2283
gt <u>c</u> Va]	aac Asn 720	Ser	att	999 Gly	aag Lys	agc Ser 725	Leu	Pro	agc Ser	gag	gcg Ala 730	Ser	gag Glu	cag Gln	tat Tyr	2331
tt <u>c</u> Leu 735	Thr	aaa Lys	gcc	tca Ser	gaa Glu 740	Pro	gat Asp	aaa Lys	aac Asn	CCC Pro	Thr	gct	gtg	gaa Glu	gga Gly 750-	2379

																•
ctg Leu	gga Gly	tca Ser	gag Glu	cct Pro 755	gat Asp	aat Asn	ttg Leu	gag Glu	att Ile 760	acg Thr	tgg Trp	aag Lys	ccc Pro	ttg Leu 765	aat Asn	2427
ggt Gly	ttc Phe	gaa Glu	tct Ser 770	aat Asn	G1 y 999	cca Pro	ggc Gly	ctt Leu 775	cag Gln	tac Tyr	aaa Lys	gtt Val	agc Ser 780	tgg Trp	cgc Arg	2475
cag Gln	aaa Lys	gat Asp 785	ggt Gly	gat Asp	gat Asp	gaa Glu	tgg Trp 790	aca Thr	tct Ser	gtg Val	gtt Val	gtg Val 795	gca Ala	aat Asn	gta Val	2523
tcc Ser	aaa Lys 800	tat Tyr	att Ile	gtc Val	tca Ser	ggc Gly 805	acg Thr	cca Pro	acc Thr	ttt Phe	gtt Val 810	cca Pro	tac Tyr	ctg Leu	atc Ile:	2571
aaa Lys 815	gtt Val	cag Gln	gcc Ala	ctg Leu	aat Asn 820	gac Asp	atg Met	ggg Gly	ttt Phe	gcc Ala 825	Pro	gag Glu	cca Pro	gct Ala	gta Val 830	2619
gtc Val	atg Met	gga Gly	cat His	tct Ser 835	gga Gly	gaa Glu	gac Asp	ctc Leu	cca Pro 840	atg Met	gtg Val	gct Ala	cct Pro	999 Gly 845	aac Asn	2667
gtg Val	cgt Arg	gtg Val	aat Asn 850	gtg Val	gtg Val	aac Asn	agt Ser	acc Thr 855	tta Leu	gcc Ala	gag Glu	gtg Val	cac His 860	tgg Trp	gac Asp	2715
cca Pro	gta Val	cct Pro 865	ctg Leu	aaa Lys	agc Ser	atc Ile	cga Arg 870	gga Gly	cac His	cta Leu	caa Gln	ggc Gly 875	tat Tyr	cgg Arg	att Ile	2763
tac Tyr	tat Tyr 880	tgg Trp	aag Lys	acc Thr	cag Gln	agt Ser 885	tca Ser	tct Ser	aa a Lys	aga Arg	aac Asn 890	aga Arg	cgt Arg	cac His	att Ile	2811
gag Glu 895	aaa Lys	aag Lys	atc Ile	ctc Leu	acc Thr 900	ttc Phe	caa Gln	ggc Gly	agc Ser	aag Lys 905	act Thr	cat His	ggc Gly	atg Met	ttg Leu 910	2859
ccg Pro	Gly 999	cta Leu	gag Glu	ccc Pro 915	Pne	agc Ser	His	Tyr	aca Thr 920	Leu	aat Asn	gtc Val	Arg	gtg Val 925	gtc Val	2907
aat Asn	999 Gly	aaa Lys	999 Gly 930	gag Glu	ggc Gly	cca Pro	gcc Ala	agc Ser 935	cct Pro	gac Asp	aga Arg	gtc Val	ttt Phe 940	aat Asn	act Thr	2955
cca Pro	gaa Glu	gga Gly 945	gtc Val	ccc Pro	agt Ser	gct Ala	ccc Pro 950	tcg Ser	tct Ser	ttg Leu	aag Lys	att Ile 955	gtg Val	aat Asn	cca Pro	3003
aca Thr	ctg Leu 960	gac Asp	tct Ser	ctc Leu	act Thr	ttg Leu 965	gaa Glu	tgg Trp	gat Asp	cca Pro	ccg Pro 970	agc Ser	cac His	ccg Pro	aat Asn	3051

ggc att ttg aca	gag tac acc	tta aag tat	cag cca att aac	agc aca 3099
Gly lie Leu Thr		Leu Lys Tyr	Gln Pro Ile Asn	
,	980		985	990
cat gaa tta ggc	cct ctg gta	gat ffg aaa	att cct gcc aac	230 202 2147
His Glu Leu Gly	Pro Leu Val	Asp Leu Lvs	Ile Pro Ala Asn	aag aca 3147
•	995	1000		005
cgg tgg act tta	aaa aat tta	aat ttc agc	act cga tat aag	ttt tat 3195
	Lys Asn Leu		Thr Arg Tyr Lys	Phe Tyr
1010		1015	1020	
ttc tat gca caa	aca tca gca	gga tca gga	agt caa att aca	ana in 2242
Phe Tyr Ala Gln	Thr Ser Ala	Glv Ser Glv	Ser Gln Ile Thr	gag gaa 3243
1025		.030	1035	014 014
gca gta aca act	gtg gat gaa	gct ggt att	ctt cca cct gat	gta ggt 3291
1040		Ala Gly Ile	Leu Pro Pro Asp	Val Gly
1040	1045		1050	
gca ggc aaa gtt	caa gct gta	aat acc acc	atc agc aat ctt	205 005 2220
Ala Gly Lys Val	Gln Ala Val	Asn Thr Arg	Ile Ser Asn Leu	act gct 3339 Thr Ala
1055	1060		1065	1070
gca gct gct gag	acc tat gcc	aat atc agt	tgg gaa tat gag	gga cca 3387
			Trp Glu Tyr Glu	
-	1075	1080	1	.085
gag cat gtg aac	ttt tat oft	gaa tat ggt	gta gca ggc agc	aaa gaa 3435
Glu His Val Asn	Phe Tyr Val	Glu Tyr Gly	Val Ala Gly Ser	Lvs Glu
1090		1095	1100	
gaa tgg aga aaa	gaa att gta	aat ggt tct	cgg agc ttc ttt	ggg tta 3483
1105		ASR GIY Ser 1110	Arg Ser Phe Phe	Gly Leu
1105	•	1110	1115	
aag ggt cta atg	cca gga aca	gca tac aaa	gtt cga gtt ggt	gct qtq 3531
Lys Gly Leu Met	Pro Gly Thr	Ala Tyr Lys	Val Arg Val Gly	Ala Val
1120	1125		1130	
000 020 tot oot	*** *** ***			
Gly Asp Ser Gly	Dhe Val Ser	Ser Clu Ner	gtg ttt gag aca Val Phe Glu Thr	ggc cca 3579
1135	1140		1145	
•			1143	1150
gcg atg gca agc	cgg cag gtg	gat att gca	act cag ggc tgg	ttc att 3627
Ala Met Ala Ser	Arg Gln Val	Asp Ile Ala	Thr Gln Gly Trp	Phe Ile
	1155	1160		1165
ggt ctg atg tot	act att	ata	***	
Gly Leu Met Cye	Ala Val Ala	Leu Iou Tia	tta att ttg ctg Leu Ile Leu Leu	att gtt 3675
1170	tat via	1175	Leu lie Leu Leu 1180	TIE VAI
tgc ttc atc aga	aga aac aag	ggt ggt aaa	tat cca gtt aaa	gaa aag 3723
Cys Phe Ile Arg	Arg Asn Lys	Gly Gly Lys	Tyr Pro Val Lys	Glu Lys
1185		1190	1195	

-	1200	, Ald		nia	wsp	1205	GIU	lle	GIn	Pro	atg Met 1210	Lys	Glu	Asp	Asp	3771
121	5	rne	GIY	GIU	1220	ser	Asp	Ala	Glu	Asp 1225	•	Lys	Pro	Leu	Lys 1230	3819
-,-	U.J	501	Arg	1235	PIO	ser	Asp	Arg	Thr 1240	Val	aaa Lys	Lys	Glu	Asp 1245	Ser	3867
			1250	val	АЗР	ıyr	GIY	G1u 1255	Gly	Val		Gly	Gln 1260	Phe	Asn	3915
	1.02	1265	Jei	FIIC	116	GIY	1270	lyr	Ser	Gly		Lys 1275	Glu	Lys	Glu	3963
	gct Ala 1280	214	gga Gly	aac Asn	Giu	agc Ser 1285	tca Ser	gag Glu	gca Ala	Pro	tct Ser 1290	cct Pro	gtc Val	aac Asn	gcc Ala	4011
atg Met 129	ASII	tcc Ser	ttt Phe	vai	taa 1300	tttt	ttaa	get (caaa	gcca	at a	tcca	attt	c		4059
tet:	gaa	tat :														
	-9-4	-90 (Lale	ccta	ag ct	ctt	gttt	gtca	agcc	ctct	cata	actat	ga a	acata	atigggt	4119
		ata (ag ct	cttg	gttt	g tca	agcc	ctct	cata	actat	iga (acata	atigggt	4119 4134
<210 <211 <212	gagt: 0> 2 l> 1: 2> P1	ata t 299	ttt	2	ag ct	ctt	gttt;	g tc:	agcc	ctct	Cata	actat	iga i	acata	atgggt	
<210 <211 <212 <213	<pre>page: 1> 2 1> 1: 2> Pi 3> Ho 1> 2</pre>	ata t 299 RT Omo s	sapie	ens												
<210 <211 <212 <213 <400 Met)> 2 l> 1: l> Pi l> Ho l> 2 Pro	ata t 299 RT Omo s	apie Lys	ens Lýs 5	Arg	Leu	Ser	Ala	Gly 10	Arg	Val	Pro	Leu	Ile	Leu	
<210 <211 <212 <213 <400 Met)> 2 l> 1: l> Pi l> Ho l> 2 Pro	ata t 299 RT Omo s	apie Lys	ens Lýs 5	Arg	Leu	Ser	Ala	Gly 10	Arg		Pro	Leu	Ile	Leu	
<210 <211 <212 <213 <400 Met 1 Phe)> 2 > 1: > 1: > P > 2 Pro	299 RT DMO s Lys Cys Glu 35	Lys Gln 20 Asp	ens Lys 5 Met Leu	Arg Ile Val	Leu Ser Gln	Ser Ala Pro 40	Ala Leu 25 Pro	Gly 10 Glu Thr	Arg Val Ile	Val Pro Thr	Pro Leu Gln 45	Leu Asp 30 Gln	Ile 15 Pro	Leu Lys Pro	
<210 213</213</400 Met 1 Phe Leu Lys</td <td>l> 2 l> 1: 2> Pi 3> Ho Pro Leu Leu Asp 50</td> <td>299 RT DMO s Lys Cys Glu 35</td> <td>Lys Gln 20 Asp</td> <td>ens Lys 5 Met Leu</td> <td>Arg Ile Val Asp</td> <td>Leu Ser Gln Pro 55</td> <td>Ser Ala Pro 40 Arg</td> <td>Ala Leu 25 Pro Glu</td> <td>Gly 10 Glu Thr</td> <td>Arg Val Ile</td> <td>Val Pro Thr Val 60</td> <td>Pro Leu Gln 45 Ile</td> <td>Leu Asp 30 Gln</td> <td>Ile 15 Pro Ser</td> <td>Leu Lys Pro Glu</td> <td></td>	l> 2 l> 1: 2> Pi 3> Ho Pro Leu Leu Asp 50	299 RT DMO s Lys Cys Glu 35	Lys Gln 20 Asp	ens Lys 5 Met Leu	Arg Ile Val Asp	Leu Ser Gln Pro 55	Ser Ala Pro 40 Arg	Ala Leu 25 Pro Glu	Gly 10 Glu Thr	Arg Val Ile	Val Pro Thr Val 60	Pro Leu Gln 45 Ile	Leu Asp 30 Gln	Ile 15 Pro Ser	Leu Lys Pro Glu	
<pre><210 <211 <212 <213 <400 Met</pre>	l> 2 l> 1: 2> Pi 3> Ho 0> 2 Pro Leu Leu Asp 50 Lys	299 RT DMO s Lys Cys Glu 35 Tyr	Lys Gln 20 Asp Ile	ens Lys 5 Met Leu Ile	Arg Ile Val Asp Pro 70	Leu Ser Gln Pro 55	Ser Ala Pro 40 Arg	Ala Leu 25 Pro Glu Phe	Gly 10 Glu Thr Asn Ser	Arg Val Ile Ile Trp 75	Val Pro Thr	Pro Leu Gln 45 Ile Arg	Leu Asp 30 Gln Gln	Ile 15 Pro Ser Cys	Leu Lys Pro Glu Thr 80	

Gly Thr Leu Ile Ile Asn Ile Met Ser Glu Gly Lys Ala Glu Thr Tyr 100 105 110

Glu Gly Val Tyr Gln Cys Thr Ala Arg Asn Glu Arg Gly Ala Ala Val

Ser Asn Asn Ile Val Val Arg Pro Ser Arg Ser Pro Leu Trp Thr Lys

Glu Lys Leu Glu Pro Ile Thr Leu Gln Ser Gly Gln Ser Leu Val Leu 145 150 155 160

Pro Cys Arg Pro Pro Ile Gly Leu Pro Pro Pro Ile Ile Phe Trp Met
165 170 175

Asp Asn Ser Phe Gln Arg Leu Pro Gln Ser Glu Arg Val Ser Gln Gly
180 185 190

Leu Asn Gly Asp Leu Tyr Phe Ser Asn Val Leu Pro Glu Asp Thr Arg 195 200 205

Glu Asp Tyr Ile Cys Tyr Ala Arg Phe Asn His Thr Gln Thr Ile Gln 210 215 220

Gln Lys Gln Pro Ile Ser Val Lys Val Ile Ser Val Asp Glu Leu Asn 225 230 235 240

Asp Thr Ile Ala Ala Asn Leu Ser Asp Thr Glu Phe Tyr Gly Ala Lys 245 250 255

Ser Ser Arg Glu Arg Pro Pro Thr Phe Leu Thr Pro Glu Gly Asn Ala 260 265 270

Ser Asn Lys Glu Glu Leu Arg Gly Asn Val Leu Ser Leu Glu Cys Ile 275 280 285

Ala Glu Gly Leu Pro Thr Pro Ile Ile Tyr Trp Ala Lys Glu Asp Gly 290 295 300

Met Leu Pro Lys Asn Arg Thr Val Tyr Lys Asn Phe Glu Lys Thr Leu 305 310 315 320

Gln Ile Ile His Val Ser Glu Ala Asp Ser Gly Asn Tyr Gln Cys Ile 325 330 335

Ala Lys Asn Ala Leu Gly Ala Ile His His Thr Ile Ser Val Arg Val 340 345 350

Lys Ala Ala Pro Tyr Trp Ile Thr Ala Pro Gln Asn Leu Val Leu Ser 355 360 365

Pro Gly Glu Asp Gly Thr Leu Ile Cys Arg Ala Asn Gly Asn Pro Lys 370 375 380

Pro-Arg-Ile-Ser-Trp-Leu-Thr-Asn-Gly-Val-Pro-Ile-Glu-Ile-Ala-Pro-385 390 395 400

Asp Asp Pro Ser Arg Lys Ile Asp Gly Asp Thr Ile Ile Phe Ser Asn 405 410 415

- Val Gln Glu Arg Ser Ser Ala Val Tyr Gln Cys Asn Ala Ser Asn Glu 420 425 430
- Tyr Gly Tyr Leu Leu Ala Asn Ala Phe Val Asn Val Leu Ala Glu Pro
 435 440 445
- Pro Arg Ile Leu Thr Pro Ala Asn Thr Leu Tyr Gln Val Ile Ala Asn 450 455 460
- Arg Pro Ala Leu Leu Asp Cys Ala Phe Phe Gly Ser Pro Leu Pro Thr 465 470 475 480
- Ile Glu Trp Phe Lys Gly Ala Lys Gly Ser Ala Leu His Glu Asp Ile 485 490 495
- Tyr Val Leu His Glu Asn Gly Thr Leu Glu Ile Lys Asp Ala Thr Trp 500 505 510
- Ile Val Lys Glu Ile Pro Val Ala Gln Lys Asp Ser Thr Gly Thr Tyr 515 520 525
- Thr Cys Val Ala Arg Asn Lys Leu Gly Met Ala Lys Asn Glu Val His 530 540
- Leu Gln Pro Glu Tyr Ala Val Val Gln Arg Gly Ser Met Val Ser Phe 545 550 555 560
- Glu Cys Lys Val Lys His Asp His Thr Leu Ser Leu Thr Val Leu Trp
 565 570 575
- Leu Lys Asp Asn Arg Glu Leu Pro Ser Asp Glu Arg Phe Thr Val Asp 580 585 590
- Lys Asp His Leu Val Val Ala Asp Val Ser Asp Asp Asp Ser Gly Thr 595 600 605
- Tyr Thr Cys Val Ala Asn Thr Thr Leu Asp Ser Val Ser Ala Ser Ala 610 615 620
- Val Leu Ser Val Val Ala Pro Thr Pro Thr Pro Ala Pro Val Tyr Asp
 625 630 635 640
- Val Pro Asn Pro Pro Phe Asp Leu Glu Leu Thr Asp Gln Leu Asp Lys 645 650 655
- Ser Val Gln Leu Ser Trp Thr Pro Gly Asp Asp Asn Asn Ser Pro Ile
 660 665 670
- Thr Lys Phe Ile Ile Glu Tyr Glu Asp Ala Met His Lys Pro Gly Leu 675 680 685
- Trp His His Gln Thr Glu Val Ser Gly Thr Gln Thr Thr Ala Gln Leu 690 695 700

Lys Leu Ser Pro Tyr Val Asn Tyr Ser Phe Arg Val Met Ala Val Asn 705 710 715 720

- Ser Ile Gly Lys Ser Leu Pro Ser Glu Ala Ser Glu Gln Tyr Leu Thr
 725 730 735
- Lys Ala Ser Glu Pro Asp Lys Asn Pro Thr Ala Val Glu Gly Leu Gly
 740 745 750
- Ser Glu Pro Asp Asn Leu Glu Ile Thr Trp Lys Pro Leu Asn Gly Phe
 755 760 765
- Glu Ser Asn Gly Pro Gly Leu Gln Tyr Lys Val Ser Trp Arg Gln Lys
 770 780
- Asp Gly Asp Asp Glu Trp Thr Ser Val Val Val Ala Asn Val Ser Lys
 785 790 795 800
- Tyr Ile Val Ser Gly Thr Pro Thr Phe Val Pro Tyr Leu Ile Lys Val 805 810 815
- Gln Ala Leu Asn Asp Met Gly Phe Ala Pro Glu Pro Ala Val Val Met 820 825 830
- Gly His Ser Gly Glu Asp Leu Pro Met Val Ala Pro Gly Asn Val Arg 835 840 845
- Val Asn Val Val Asn Ser Thr Leu Ala Glu Val His Trp Asp Pro Val 850 860
- Pro Leu Lys Ser Ile Arg Gly His Leu Gln Gly Tyr Arg Ile Tyr Tyr 865 870 875 880
- Trp Lys Thr Gln Ser Ser Ser Lys Arg Asn Arg Arg His Ile Glu Lys 885 890 895
- Lys Ile Leu Thr Phe Gln Gly Ser Lys Thr His Gly Met Leu Pro Gly 900 905 910
- Leu Glu Pro Phe Ser His Tyr Thr Leu Asn Val Arg Val Val Asn Gly
 915 920 925
- Lys Gly Glu Gly Pro Ala Ser Pro Asp Arg Val Phe Asn Thr Pro Glu 930 935 940
- Gly Val Pro Ser Ala Pro Ser Ser Leu Lys Ile Val Asn Pro Thr Leu 945 950 955 960
- Asp Ser Leu Thr Leu Glu Trp Asp Pro Pro Ser His Pro Asn Gly Ile 965 970 975
- Leu Thr Glu Tyr Thr Leu Lys Tyr Gln Pro Ile Asn Ser Thr His Glu 980 985 990
- Leu-Gly-Pro-Leu-Val-Asp-Leu-Lys-Ile-Pro-Ala-Asn-Lys-Thr-Arg-Trp-995 1000 1005

Thr Leu Lys Asn Leu Asn Phe Ser Thr Arg Tyr Lys Phe Tyr 1010 1015 1020

- Ala Gln Thr Ser Ala Gly Ser Gly Ser Gln Ile Thr Glu Glu Ala Val 025 1030 1035 1040
- Thr Thr Val Asp Glu Ala Gly Ile Leu Pro Pro Asp Val Gly Ala Gly
 1045 1050 1055
- Lys Val Gln Ala Val Asn Thr Arg Ile Ser Asn Leu Thr Ala Ala Ala 1060 1065 1070
- Ala Glu Thr Tyr Ala Asn Ile Ser Trp Glu Tyr Glu Gly Pro Glu His 1075 1080 1085
- Val Asn Phe Tyr Val Glu Tyr Gly Val Ala Gly Ser Lys Glu Glu Trp 1090 1095 1100
- Arg Lys Glu Ile Val Asn Gly Ser Arg Ser Phe Phe Gly Leu Lys Gly 105 1110 1115 1120
- Leu Met Pro Gly Thr Ala Tyr Lys Val Arg Val Gly Ala Val Gly Asp 1125 1130 1135
- Ser Gly Phe Val Ser Ser Glu Asp Val Phe Glu Thr Gly Pro Ala Met 1140 1145 1150
- Ala Ser Arg Gln Val Asp Ile Ala Thr Gln Gly Trp Phe Ile Gly Leu 1155 1160 1165
- Met Cys Ala Val Ala Leu Leu Ile Leu Ile Leu Leu Ile Val Cys Phe 1170 1175 1180
- Ile Arg Arg Asn Lys Gly Gly Lys Tyr Pro Val Lys Glu Lys Glu Asp 185 1190 1195 1200
- Ala His Ala Asp Pro Glu Ile Gln Pro Met Lys Glu Asp Asp Gly Thr 1205 1210 1215
- Phe Gly Glu Tyr Ser Asp Ala Glu Asp His Lys Pro Leu Lys Lys Gly
 1220 1225 1230
- Ser Arg Thr Pro Ser Asp Arg Thr Val Lys Lys Glu Asp Ser Asp Asp 1235 1240 1245
- Ser Leu Val Asp Tyr Gly Glu Gly Val Asn Gly Gln Phe Asn Glu Asp 1250 1255 1260
- Gly Ser Phe Ile Gly Gln Tyr Ser Gly Lys Lys Glu Lys Glu Pro Ala 265 1270 1275 1280
- Glu Gly Asn Glu Ser Ser Glu Ala Pro Ser Pro Val Asn Ala Met Asn 1285 1290 1295

Ser Phe Val

<210> 3

· •

```
<211> 38
<212> DNA
<213> Homo sapiens
<400> 3
tctcatacta tgaacatatg ggtagagagt atattttc
                                                                   38
<210> 4
<211> 123
<212> DNA
<213> Rattus norvegicus
<400> 4
tctcatacta tggacatatg ggtagaaaga atgttttctg cggtatatga gtattataag 60
aacagagcaa gaacataact cagtcagtca gatgatacgt taatatgaac tggggtgaaa 120
agg
                                                                   123
<210> 5
<211> 176
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: clone D4-1
tctcatacta tgaacatatg ggtagagagt atattttctg ctgtatgtta gtattatgag 60
aatagttaca gcaaaaacat aactcagtca aagtatatgt taatatgaac tggaatgcaa 120
aagtgcatac tttttcattc aaaatgggta ttcttgattt cctaaaaaaa aaaaaa
                                                                    176
<210> 6
<211> 38
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: primer
tagatacaac tagtcaatgc ctctaatgaa tatggata
                                                                    38
<210> 7
<211> 38
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: primer
```



<400> 7

WO 99/55380	PCT/US99/09039
agatagatee geggaatagt aaateegata geettgta	38
<210> 8 <211> 15 <212> DNA <213> Artificial Sequence	
<220> <221> modified_base <222> (1) <223> n=a, c, g, or t	
<220> <223> Description of Artificial Sequence: primer	e e
<400> B ngctgctctc atact	15
<210> 9 <211> 24 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence: primer	
<400> 9 aacatatggg tagagagtat attt	24
<210> 10 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence: primer	
<400> 10 ctttgcattc cagttcatat taa	23
<210> 11 <211> 20 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence: primer	
<400> 11 tgtggtgaca gatcacggct	20
<210> 12	

```
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: primer
<400> 12
cagcicaaac ctgtgatttc c
                                                                    21
<210> 13
<211> 60
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: primer
<400> 13
aataggtatt ggtgaattta aagactcact ctccataaat gctacgaata ttaaacactt 60
<210> 14
<211> 21
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer
<400> 14
cggagcaata tgaaatgatc t
                                                                    21
<210> 15
<211> 19
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer
<400> 15
gcaaatacag ctcctattg
                                                                    19
<210> 16
<211> 43
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer
<400> 16
gctgtatgtt agtattatga gaatagttac agcaaaaaca taa
```

PCT/US99/09039

43

WO 99/55380

WO 99/55380	PCT/US99/09039
<210> 17 <211> 40	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: primer	
<400> 17	
taggeetgae tggeattgta ttageaaact cateaetaga	40
<210> 18	·
<211> 37	
<212> DNA	
<213> Artificial Sequence	
<220> <223> Description of Artificial Sequence: primer	
<400> 18	
tagatacaac tagtctaatg cagcttaaaa taatgcc	
	37
<210> 19	
<211> 39	
<212> DNA <213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: primer	
<400> 19	
agatagatcc gcggatatcc atattcatta gaggcattg	39
<210> 20	
<211> 38	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: primer	
<400> 20	
tagatacaac tagtcaatgc ctctaatgaa tatggata	38
<210> 21	
<211> 38	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: primer	
<400> 21	
agatagatcc gcggaatagt aaatccgata gccttgta	38

```
<210> 22
<211> 61
<212> DNA
<213> Homo sapiens
<220>
<223> Description of Artificial Sequence: primer
<400> 22
aggagttaag atgctaatgc agcttaaaat aatgccgaaa aagaagcgct tatctgcggg 60
                                                                    61
<210> 23
<211> 19
<212> DNA
<213> Homo sapiens
<400> 23
cattagcatc ttaactcct
                                                                    19
<210> 24
<211> 21
<212> DNA
<213> Homo sapiens
<400> 24
tcggcattat tttaagctgc a
                                                                     21
<210> 25
<211> 17
<212> DNA
<213> Homo sapiens
<400> 25
gcagataagc gcttctt
                                                                     17
<210> 26
<211> 20
<212> DNA
<213> Homo sapiens
<400> 26
actagagata cagatcatat
                                                                     20
 <210> 27
 <211> 20
 <212> DNA
 <213> Homo_sapiens_
 <400> 27
 catatacgat cgatcgatgc
                                                                     20
```

```
<210> 28
<211> 20
<212> DNA
<213> Homo sapiens
<400> 28
gatagtgctg atcgatgcta
                                                                   20
<210> 29
<211> 48
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer
<400> 29
catacgaatt ctagatacaa ctagtctaat gcagcttaaa ataatgcc
                                                                   4 B
<210> 30
<211> 50
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer
<400> 30
agatagatee geggatatee atatteatta gaggeattgg gateeeatae
                                                                   50
<210> 31
<211> 1371
<212> DNA
<213> Homo sapiens
<400> 31
atgeegaaaa agaagegett atetgeggge agagtgeece tgattetett eetgigeeag 60
atgattagtg cactggaagt acctcttgat ccaaaacttc ttgaagactt ggtacagcct 120
ccaaccatca cccaacagtc tccaaaagat tacattattg accctcggga gaatattgta 180
atccagtgtg aagccaaagg gaaaccgccc ccaagctttt cctggacccg taatgggact 240
cattttgaca tcgataaaga ccctctggtc accatgaagc ctggcacagg aacgctcata 300
attaacatca tgagcgaagg gaaagctgag acctatgaag gagtctatca gtgtacagca 360
aggaacgaac gcggagctgc agtttctaat aacattgttg tccgcccatc cagatcacca 420
ttgtggacca aagaaaaact tgaaccaatc acacttcaaa gtggtcagtc tttagtactt 480
ccctgcagac ccccaattgg attaccacca cctataatat tttggatgga taattccttt 540
```

<400> 32

<210> 32

<211> 1371

<212> DNA

<213> Rattus norvegicus

gacactatag	ctgctaattt	gagtgacact	gagttttatg	gtgctaaatc	tagtaaagag	780
aggccaccaa	catttctaac	tccagagggc	aatgaaagtc	acaaggaaga	attaagagga	840
aacgtgcttt	ccctggagtg	cattgcagaa	ggcctaccta	ctccagttat	ttactggatc	900
aaggaagatg	gaacgcttcc	tgtcaaccgg	acgttttatc	ggaactttaa	gaaaaccttg	960
cagatcattc	atgtctctga	agcagactct	ggaaattatc	agtgcatagc	aaaaaacgca	1020
ttgggagccg	tccatcatac	catttctgtc	acagttaaag	cggctcccta	ctggattgtt	1080
gcacctcaca	acctcgtgct	ttccccaggg	gagaatggga	ccctcatctg	cagagetaac	1140
ggcaacccaa	aacccagaat	tagctggtta	acaaatggag	tcccagtaga	aattgctctc	1200
gatgacccca	gccgaaaaat	cgatggtgat	accattatgt	tttcaaatgt	tcaagaaagc	1260
tcaagtgcgg	tttatcagtg	caatgcctct	aacaaatatg	gatatttact	agcaaatgca	1320
tttgtaaatg	tgctcgctga	accacctcgg	attcttacct	cagcaaacac	a	1371

INTERNATIONAL SEARCH REPORT

International application No. PCT/US99/09039

			
IPC(6) :	SSIFICATION OF SUBJECT MATTER A61K 48/00; CO7H 21/04 514/44; 536/24.5		
According to	o International Patent Classification (IPC) or to both n	national classification and IPC	
	DS SEARCHED		
Minimum de	ocumentation searched (classification system followed	by classification symbols)	
	514/44; 536/24.5		
Documentat	ion searched other than minimum documentation to the e	extent that such documents are included	in the fields searched
Electronic d	ata base consulted during the international search (nar	ne of data base and, where practicable	e, search terms used)
	N CAPLUS ms: nr-cam, neural-glial-related or associated-cellular	adhesion molecule, antisense	
c. Doc	UMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where app	propriate, of the relevant passages	Relevant to claim No.
Y, P	US 5,766,888 A (SOBOL et al) 16 Jun	ne 1998, col. 45-46.	15-16 20-21
Y,P	US 5,872,225 A (LEMMON) 16 Febracol. 1-2.	ruary 1999,	1-21
A,P	US 5,789,573 A (BAKER et al) 04 A 64.	august 1998, col 1, lines 41-	1-21
A	STANLEY CROOKE, Basic Principle Antisense Research and Application Heidelberg, 07 July 1998, p. 2-3.		1-21
Furt	her documents are listed in the continuation of Box C	. See patent family annex.	
• 5	pecial categories of cited documents:	"T" later document published after the ir date and not in conflict with the sp	ternational filing date or priority
	ocument defining the general state of the art which is not considered be of particular relevance	the principle or theory underlying the	he invention
.E. e	arlier document published on or after the international filing date	"X" document of particular relevance; considered novel or cannot be considered when the document is taken alone	the claimed invention eannot be dered to involve an inventive step
ci	ocument which may throw doubts on priority claim(s) or which is ited to establish the publication date of another citation or other social reason (as specified)	"Y" document of particular relevance;	the claimed invention cannot be
·0• d	ocument referring to an oral disclosure, use, exhibition or other acans	considered to involve an inventi- combined with one or more other su being obvious to a person skilled in	ich documents, such combination
	ocument published prior to the international filing date but later than	*&* document member of the same pate	ent family
	e actual completion of the international search	Date of mailing of the international s	earch report
06 JULY		/25 AU	61999
Commissi Box PCT	mailing address of the ISA/US ioner of Patents and Trademarks on, D.C. 20231	Authorized officer Authorized officer Authorized officer	(allest of "
1	No. (703) 305-3230	Telephone No. / (703) 308-0196	<i>k</i>